

The Effect of Phytoadditive Combination and Growth Performances and Meat Quality in Rainbow Trout (*Oncorhynchus mykiss*)

Erol-Florian Gabor¹, Aurel Șara¹, Mihai Bențea¹, Călina Creța², Anca Baciuc³

¹University of Agricultural Sciences and Veterinary Medicine Cluj-Napoca,
400372-Cluj-Napoca, Manastur, 3-5, Romania

²National Institute of Public Health, Teritorial Center of Public Health Cluj,
400349-Cluj-Napoca, Pasteur, 6, Romania

³SC ProPlanta SRL, Cluj Napoca, 400372-Cluj-Napoca, Mănăștur, 5-7, Romania

Abstract

Phytoadditives have been used for millennia in traditional human medicine but only recent in animals to improve the growth performances and to reduce disease outbreaks. The goal of the present experiment has been to determine the effect of phytoadditive combinations on growth performances and meat quality in rainbow trout (*Oncorhynchus mykiss*). The phytoadditive combinations were represented by garlic (2%) + ginger (1%) – group 1E and oregano (1%) and Echinacea (0.5%) – group 2E. the dietary treatment with phytoadditive combinations led to a better body weight gain in all the experimental groups compared to the control. Also the specific growth rate and the FCR values were improved in the experimental groups, the best values being recorded in group 1E. the experimental groups also had a higher survival rate, the highest value being recorded in group 1E. meat quality was not significantly influenced by dietary treatment of phytoadditive combinations. The meat Selenium content significantly varied among all experimental groups, the highest value being recorded in group 1E and the lowest in the control group. Dietary treatment with phytoadditive combination had a positive influence on growth and consumption indices with no significant changes in meat quality but significantly influencing the meat Selenium content.

Keywords: Echinacea, garlic, ginger, oregano, selenium, trout

1. Introduction

Phytoadditives are fodder additives obtained from plants or plant extracts. Their beneficial effects in humans are known for millennia and recently they have been used on animals in order to enhance the productive performances and/or to control and fight disease.

A member of the *Liliaceae* family, garlic (*Allium sativum*) has been used for centuries as a spice but also in traditional medicine. Garlic is originating in Asia Minor and spontaneously grows in southern Europe, but in cultures can be found all

over the world. It is a rich source of calcium, phosphorus and vitamin B₁; it has a high content of carbohydrates and as a consequence a high nutritive value. Garlic also contains the vitamin complex B, vitamins A, C and F [1]. Garlic is hypo-lipidemic [2], antimicrobial [3], anti-hypertensive [4], has hepato-protective and insecticidal effects [5] and also anti-fungal effects [6]. Also garlic reduces the cholesterol levels in blood serum [7, 8] by suppressing the lipid synthesis in liver [9] and increases the clotting time of blood [7]. S-allyl cysteine found in crushed garlic has inhibitory effects in tumor metabolism and enhances the immune response [2]. Allyl-sulphates enhance the S-transferase enzyme system, which plays a major role in liver

*Corresponding author; Gabor Erol, 0747040538,
Email: gogul3us@yahoo.com

detoxification from carcinogen substances. Garlic has immunostimulating activities that include the stimulation of lymphocyte synthesis, cytokine releases and stimulation of the phagocytosis [10]. Ginger is a root consumed as a delicacy, medicine or spice. Ginger is the rhizome of the *Zingiber officinale* plant, from *Zingiberaceae* family. Other notable members of this botanical family are turmeric (curcuma), cardamom and the galangal (blue ginger).

The medicinal form of ginger was called *Jamaica ginger*; it's considered as stimulant and carminative (efficient against flatulence) and was frequently used to treat indigestion, gastro paresis, constipation and cramps. The ginger lowers cholesterolemia and thins the blood, being indicated in the treatment of heart diseases; ginger also has anti-inflammatory effect. Recent researches suggest that ginger can be used in diabetes treatment [11,12]. Ginger also has analgesic, sedative, antipyretic (reduces fever) and anti-bacterial effects. In laboratory trials, ginger reduced the occurrence of skin cancer in mice and destroyed the ovary cancer cells [13,14,15].

Oregano is a species of the *Origanum* genus, *Lamiaceae* family. Spontaneously, it can grow in the temperate-warm regions of western and south-western Eurasia and in the Mediterranean region. Oregano has a strong antimicrobial and antioxidant action due to its high content of phenolic acids and flavonoids. The main chemical constituents are carvacrol, thymol, limonene, pinene, ocimene, caryophyllene. The leaves and flowers have a mild tonic effect. Oregano has a broad spectrum of use: antiseptic, antispasmodic, carminative, colagogue, emmenagogue and expectorant. Aqueous extracts, capsules or oil extracts are orally ingested in the treatment for cold, flu, mild fevers, fungal infections, indigestion and intestinal parasites. Also, oregano has a strong sedative effect, the use of high doses being counter indicated. Due to its high content of thymol, oregano is considered a very good antiseptic.

Echinacea is a genus comprised from 9 species of plants, belonging to the *Asteraceae* family. The plants have big composite flowers that bloom from the end of spring until the end of summer. From this genus, only 3 are used as medicinal plants: *Echinacea angustifolia*, *Echinacea pallida* and *Echinacea purpurea*. Echinacea extracts have immunostimulatory and anti-inflammatory

properties, for this reason the different Echinacea species take up an important place in the medicinal plant products market.

2. Materials and methods

The research has been carried out in the ICAS farm from Gilau, Cluj County on a total number of 600 rainbow trout (*Oncorhynchus mykiss*) distributed in 3 groups (one control group and 2 experimental groups) consisting of 200 fishes/group. The mean initial weight at the beginning of the experiment was 91.90g. The fish were reared in concrete tanks (L/W/H = 4/1/1.5), in the same rearing conditions (water quality, rearing density, feeding regimes, fodder quantity/quality). The experimental period was 95 days (25.06.2010 – 27.09.2010). the additives were purchased from the local market, dried and grounded to a fine powder. The fodder used was purchased from Skretting (41% crude protein, 12% crude fat, 7.8% ash). The fodder was supplemented with the additives combinations as follows: control group – base diet, group 1E – garlic (2%) + ginger (1%) and group 2E – oregano (1%) + Echinacea (0.5%). During the experimental period the body weight evolution, the weight gain, the growth rate, the feed conversion ratio (FCR) and the survival rate were monitored. At the end of the experimental period meat samples were collected for determination of crude chemical composition of meat and meat Selenium content. The meat Selenium content was carried out using the Atomic Absorption Spectroscopy method. The data collected were statistically analyzed by means of ANOVA (Tukey test). Data are expressed as mean \pm standard error of mean. All analyses were made using the GraphPad Software, (v3.05).

3. Results and discussion

Dietary use of phytoadditives combinations had a beneficial effect on body weight gain (Table 1), the highest value being recorded in group 1E (247.08g) and the lowest in the control group (227.1g). The body mass evolution was constant throughout the experimental period, with no regressions, indicating that there were no stressing

factors such as manipulations or change in feed (at the beginning of the experiment). The specific growth rate was also enhanced in all the experimental groups, the highest value being recorded in group 1E (2.60 g/day) and the lowest in the control group (2.39 g/day). FCR recorded in the experimental groups was lower than the FCR recorded in the control group; the lowest value was recorded in group 1E (1.455:1), followed by

group 2E (1.472:1); the control group presented the highest FCR value (1.518:1). These results can be explained by the various effects of the additives used: garlic enhances the performances of the intestinal microbiota [16] while oregano and ginger stimulate the appetite [17,18]. Enhancements of the growth and consumption parameters were also reported by Metwally [19]

Table 1. Growth and consumption indices recorded throughout the experimental period

Issue	C	L1E	L2E
		Garlic+ginger	Oregano+Echinacea
Initial weight (25.06.2011)	91.90	91.90	91.90
July (26.07.2011)	169.20	152.44	161.93
August (27.08.2011)	239.42	256.53	245.94
Final weight (27.09.2011)	319.00	338.98	323.98
Weight gain (g)	227.1	247.08	232.08
Specific growth rate (g/day)	2.39	2.60	2.44
FCR	1.518	1.455	1.472

The combined use of phytoadditives led to the improvement of the survival rates and subsequently to reduced losses in both the experimental groups compared to the control group (Table 2). The highest survival rates were recorded in group 1E (92.5%) followed by group 2E (87.5%); the control group had the lowest

survival rate (86.5%). This fact can be explained by the content of the phytoadditives in various active substances (allicin, zingerone, thymol, carvacrol, phenols, etc.), these results confirming the data reported by Salah (2008) [20], Metwally (2009) [19] and Zheng *et al.* (2009) [21].

Table 2. Survival rate and losses recorded during the experimental period

Issue	C	L1E	L2E
		Garlic+ginger	Oregano+Echinacea
Initial number	200	200	200
Final number	173	185	175
Losses	27	15	25
Survival (%)	86.5	92.5	87.5

Regarding the crude chemical composition of meat (table 3) an non-significant increase in dry matter (DM) content can be observed in the experimental groups (25.43 ± 0.39 – L1E, 25.43 ± 0.23 – L2E) compared to the control group (24.98 ± 0.29). The increase of body water content was also reported by Shalaby *et al.* (2006) and Metwally (2009) after dietary treatment with garlic and by Zheng *et al.*, (2009) after dietary treatment using oregano essential oil. The crude protein (CP) content varied non-significantly, lower values being recorded in both experimental groups compared to the control group

($19.76 \pm 0.34\%$). Shalaby *et al.* (2006) [22] and Metwally (2009) [19] reported an increase in the protein content after dietary application of garlic; Zheng *et al.* (2009) [21] reported the same trend using oregano essential oil. The crude fat (CF) content varied non-significantly, the values recorded in the experimental groups ($5.14 \pm 0.28\%$ – L1E, $5.47 \pm 0.31\%$ – L2E) being higher than the control group ($4.85 \pm 0.23\%$). A decrease in the crude fat content was also reported by Shalaby *et al.* (2006) [22] and Metwally (2009) [19] after dietary application of garlic on African catfish (*Clarias lazera*) and Nile tilapia (*Oreochromis*

niloticus) and by Zheng et al., (2009) [21] after administering oregano essential oil on channel catfish (*Ictalurus punctatus*). Al-Salahy (2002) [23], after dietary application of onion and garlic juices in African catfish, reported an increase of the crude protein content in meat, although the

levels of plasma total lipids lowered. The ash content varied non-significantly among the treatment groups, the highest value being recorded in group 2E followed by the control group; group 1E had the lowest value of ash content among all the treatment groups.

Table 3. Chemical composition of rainbow trout meat

Issue	M	1E	2E
		Garlic+ginger	Oregano+Echinacea
DM (%)	24.98 ± 0.29	25.43 ± 0.39	25.43 ± 0.23
CP (%)	19.76 ± 0.34	19.22 ± 0.28	19.50 ± 0.22
CF (%)	4.85 ± 0.23	5.14 ± 0.28	5.47 ± 0.31
Ash (%)	1.58 ± 0.03	1.56 ± 0.02	1.64 ± 0.11

The Selenium content of meat varied non-significantly among the treatment groups, the highest value being recorded in group 1E (176.68±8.47 µg/kg) followed by group 2E (140.06±3.17 µg/kg); the lowest value was

recorded in the control group (112.66±3.24 µg/kg). These results can be explained by the amount of Selenium found in each additive (garlic–0.4µg/g, ginger–0.7µg/100g, oregano–0.1µg/g)

Table 4. Selenium content of meat at the end of the experimental period

Issue	M	1E	2E
		Garlic+ginger	Oregano+Echinacea
Se µg/kg	112.66 ± 3.24	176.68 ± 8.47	140.06 ± 3.17
Table of significance			
	Difference of means	q	p
M vs 1E	-64.020	11.540	P<0.001 ***
M vs 2E	-27.400	4.939	P<0.05 *
1E vs 2E	36.620	6.601	P<0.01 **

4. Conclusions

The combined use of phytoadditives had a positive effect on the growth and consumption indices, leading to increased body weight gain, a superior growth rate and a lower FCR; the survival rate was also improved by the phytoadditives treatment.

The meat quality was non-significantly influenced by the use of phytoadditives combinations by increasing the dry matter content and the crude fat content and the decreasing the crude protein content of meat.

The meat Selenium content varied significantly in all treatment groups, highlighting the contribution of these phytoadditives in Selenium assimilation; these additives can be a source of Selenium, setting up the basis for a functional food.

References

1. Drăgan, S., Gergen, I., Socaciu, C. (2008) – Alimentația funcțională cu componente bioactive naturale în sindromul metabolic; Ed. Eurostampa, Timișoara, p. 200-202, 160-161, 314
2. Sumiyoshi, H., New pharmacological activities of garlic and its constituents (Review), *Folia Pharmacological Japonica*, 1997, 110, Suppl, 1, pp. 93 – 97.
3. Kumar, M. and Berwal, J. S., Sensitivity of food pathogens to garlic (*Allium sativum* L.). *J. Appl. Microbiol.*, 1998, 84, 213–215.
4. Suetsuna, K., Isolation and characterization of angiotensin I converting enzyme inhibitor dipeptides derived from *Allium sativum* (garlic). *J. Nutr. Biochem.*, 1998, 9, 415–419
5. Wang, B. H., K. A. Zuel, K. Rahaman and D. Billington., Protective effects of aged garlic extract against bromobenzene toxicity to precision cut rat liver slices. *Toxicology*, 1998, 126, 13–222.

6. Fromthing, R. A. and G. S. Bulmer, In vitro effect of aqueous extract of garlic (*Allium sativum*) on the growth and viability of *Cryptococcus neoformans*. *Mycologia*, 1978, 70, 397–405.
7. Bordia A., H. C. Bansal, S. K. Arora and S. V. Singh Effect of essential oils of garlic and onion on alimentary hyperlipemia. *Atherosclerosis*, 1975, 21, 15–19.
8. Augusti K. T. Hypocholesterolaemic effect of garlic, *Allium sativum*, Linn. *Indian J. Exp. Biol.*, 1977, 15, 89–490
9. Chang. M.W. and Johnson, M. Effect of garlic on carbohydrate metabolism and lipid synthesis in rats. *J. Nutr.*, 1980, 110, 931–936
10. Kyo. E., N. Uda, A. Suzuki, M. Kakimoto, M. Ushijima, S. Kasuga and Y. Itakura Immunomodulation and antitumor activities of aged garlic extract. *Phytomedicine*, 1998, 5, 259–267.
11. Al-Amin, Zainab M. et al.; Thomson, M; Al-Qattan, KK; Peltonen-Shalaby, R; Ali, M Anti-diabetic and hypolipidaemic properties of ginger (*Zingiber officinale*) in streptozotocin-induced diabetic rats; *British Journal of Nutrition* (Cambridge University Press), 2006, 96.
12. Afshari, Ali Taghizadeh et al.; Shirpoor, A; Farshid, A; Saadatian, R; Rasmi, Y; Saboory, E; Ilkhanizadeh, B; Allameh, A "The effect of ginger on diabetic nephropathy, plasma antioxidant capacity and lipid peroxidation in rats". *Food Chemistry* (Elsevier) 2007, 101 (1), 148–153
13. Rhode, J.; Fogoros, S.; Zick, S.; Wahl, H.; Griffith, K. A.; Huang, J.; Liu, J. R., "Ginger inhibits cell growth and modulates angiogenic factors in ovarian cancer cells". *BMC Complementary & Alternative Medicine*, 2007, 7, 44.
14. Kim, J. S.; et al. "Cytotoxic components from the dried rhizomes of *Zingiber officinale* Roscoe". *Archives of Pharmacal Research* 2008, 31 (4), 415–418.
15. Choudhury, D.; et al. "Aqueous extract of ginger shows antiproliferative activity through disruption of microtubule network of cancer cells". *Food and Chemical Toxicology*, 2010, 48 (10), 2872–2880
16. Khalil, R. H., B. M. Nadia and M. K. Soliman Effects of Biogen and Levamisol Hcl on the immune response of cultured *Oreochromis niloticus* to *Aeromonas hydrophila* vaccine. *Beni-Suef Vet. Med. J.*, Egypt, 2001, XI (2), 381-392.
17. Helene Christine Reinbach*, Torben Martinussen, Per Møller - Effects of hot spices on energy intake, appetite and sensory specific desires in humans, *Food Quality and Preference*, 2010, 21, 655–661
18. Yeomans MR, Gray RW, Mitchell CJ, True S. Independent effects of palatability and within-meal pauses on intake and appetite ratings in human volunteers. *Appetite*. 1997, 29(1), 61-76.
19. Metwally M.A.A. - Effects of Garlic (*Allium sativum*) on Some Antioxidant Activities in Tilapia Nilotica (*Oreochromis niloticus*); *World Journal of Fish and Marine Sciences*, 2009, 1 (1), 56-64
20. Salah M. A., Mohamed F. M., George J. - Echinacea as immunostimulatory agent in Nile tilapia (*Oreochromis niloticus*) via earthen ponds experiment; 8th International Symposium on Tilapia in Aquaculture 2008.
21. Zheng Z.L., Tan J. Y.W., Liu H.Y., Zhou X.H., Xiang X., Wang K.Y. - Evaluation of oregano essential oil (*Origanum heracleoticum* L.) on growth, antioxidant effect and resistance against *Aeromonas hydrophila* in channel catfish (*Ictalurus punctatus*); *Aquaculture*, 2009, 292, 214–218
22. Shalaby A. M., Khattab Y. A., Abdel R. A. M. – Effects of garlic (*Allium sativum*) and chloramphenicol on growth performance, physiological parameters and survival of Nile tilapia (*Oreochromis niloticus*); *J. Venom. Anim. Toxins incl. Trop. Dis.*, 2006, 12, 2, 172-201
23. Al-Salahy M.B. - Some physiological studies on the effect of onion and garlic juices on the fish, *Clarias lazera*; *Fish Physiology and Biochemistry*, 2002, 27: 129–142,.