

Supplement to *Critical Care Alert* September 2021; Volume 29, Number 6

REFERENCES

- Star RA. Treatment of acute renal failure. *Kidney Int* 1998;54:1817-1831.
- Sharma A, Mucino MJ, Ronco C. Renal functional reserve and renal recovery after acute kidney injury. *Nephron Clin Pract* 2014;127:94-100.
- Musso CG, Michelangelo H, Vilas M, et al. Creatinine reabsorption by the aged kidney. *Int Urol Nephrol* 2009;41:727-731.
- Doi K, Yuen PST, Eisner C, et al. Reduced production of creatinine limits its use as marker of kidney injury in sepsis. *J Am Soc Nephrol* 2009;20:1217-1221.
- Siew ED, Peterson JF, Eden SK, et al. Use of multiple imputation method to improve estimation of missing baseline serum creatinine in acute kidney injury research. *Clin J Am Soc Nephrol* 2013;8:10-18.
- Pickkers P, Darmon M, Hoste E, et al. Acute kidney injury in the critically ill: An updated review on pathophysiology and management. *Intensive Care Med* 2021;47:835-850.
- Seguro AC, Lomar AV, Rocha AS. Acute renal failure of leptospirosis: Nonoliguric and hypokalemic forms. *Nephron* 1990;55:146-151.
- Kulvichit W, Kellum JA, Srisawat N. Biomarkers in acute kidney injury. *Crit Care Clin* 2021;37:385-398.
- Rizvi MS, Kashani KB. Biomarkers for early detection of acute kidney injury. *J Appl Lab Med* 2017;2:386.
- Chawla LS, Davison DL, Brasha-Mitchell E, et al. Development and standardization of furosemide stress test to predict the severity of acute kidney injury. *Crit Care* 2013;17:R207.
- Ho J, Tangri N, Komenda P, et al. Urinary, plasma, and serum biomarkers' utility for predicting acute kidney injury associated with cardiac surgery in adults: A meta-analysis. *Am J Kidney Dis* 2015;66:993-1005.
- Nickolas TL, Schmidt-Ott KM, Canetta P, et al. Diagnostic and prognostic stratification in the emergency department using urinary biomarkers of nephron damage: A multicenter prospective cohort study. *J Am Coll Cardiol* 2012;59:246-255.
- Haase M, Bellomo R, Devarajan P, et al. Accuracy of neutrophil gelatinase-associated lipocalin (NGAL) in diagnosis and prognosis in acute kidney injury: A systematic review and meta-analysis. *Am J Kidney Dis* 2009;54:1012-1024.
- Coll E, Botey A, Alvarez L, et al. Serum cystatin C as a new marker for noninvasive estimation of glomerular filtration rate and as a marker for early renal impairment. *Am J Kidney Dis* 2000;36:29-34.
- Herget-Rosenthal S, Marggraf G, Hüsing J, et al. Early detection of acute renal failure by serum cystatin C. *Kidney Int* 2004;66:1115-1122.
- Nejat M, Pickering JW, Walker RJ, Endre ZH. Rapid detection of acute kidney injury by plasma cystatin C in the intensive care unit. *Nephrol Dial Transplant* 2010;25:3283.
- Kashani K, Al-Khafaji A, Ardiles T, et al. Discovery and validation of cell cycle arrest biomarkers in human acute kidney injury. *Crit Care* 2013;17:R25.
- Singer E, Elger A, Elitok S, et al. Urinary neutrophil gelatinase-associated lipocalin distinguishes pre-renal from intrinsic renal failure and predicts outcomes. *Kidney Int* 2011;80:405.
- Tecson KM, Erhardtson E, Eriksen PM, et al. Optimal cut points of plasma and urine neutrophil gelatinase-associated lipocalin for the prediction of acute kidney injury among critically ill adults: Retrospective determination and clinical validation of a prospective multicentre study. *BMJ Open* 2017;7:e016028.
- Huelin P, Sola E, Elia C, et al. Neutrophil gelatinase-associated lipocalin for assessment of acute kidney injury in cirrhosis: A prospective study. *Hepatology* 2019;70:319-333.
- Koyner JL, Shaw AD, Chawla LS, et al. Tissue inhibitor metalloproteinase-2 (TIMP-2)-IGF-binding protein-7 (IGFBP7) levels are associated with adverse long-term outcomes in patients with AKI. *J Am Soc Nephrol* 2015;26:1747-1754.
- Lumlertgul N, Peerapornratana S, Trakamvanich T, et al. Early versus standard initiation of renal replacement therapy in furosemide stress test non-responsive acute kidney injury patients (the FST trial). *Crit Care* 2018;22:101.
- Srisawat N, Murugan R, Lee M, et al. Plasma neutrophil gelatinase-associated lipocalin predicts recovery from acute kidney injury following community-acquired pneumonia. *Kidney Int* 2011;80:545-552.
- Srisawat N, Wen X, Lee M, et al. Urinary biomarkers and renal recovery in critically ill patients with renal support. *Clin J Am Soc Nephrol* 2011;6:1815-1823.
- Koyner JL, Bennett MR, Worcester EM, et al. Urinary cystatin C as an early biomarker of acute kidney injury following adult cardiothoracic surgery. *Kidney Int* 2008;74:1059-1069.
- Spahillari A, Parikh CR, Sint K, et al. Serum cystatin C- versus creatinine-based definitions of acute kidney injury following cardiac surgery: A prospective cohort study. *Am J Kidney Dis* 2012;60:922-929.
- Xie Y, Ankawi G, Yang B, et al. Tissue inhibitor metalloproteinase-2 (TIMP-2) • IGF-binding protein-7 (IGFBP7) levels are associated with adverse outcomes in patients in the intensive care unit with acute kidney injury. *Kidney Int* 2019;95:1486-1493.
- Meersch M, Schmidt C, Hoffmeier A, et al. Prevention of cardiac surgery-associated AKI by implementing the KDIGO guidelines in high-risk patients identified by biomarkers: The PrevAKI randomized controlled trial. *Intensive Care Med* 2017;43:1551-1561.
- Hoste E, Bihorac A, Al-Khafaji A, et al. Identification and validation of biomarkers of persistent acute kidney injury: The RUBY study. *Intensive Care Med* 2020;46:943-953.
- Liangos O, Perianayagam MC, Vaidya VS, et al. Urinary N-acetyl-beta-(D)-glucosaminidase activity and kidney injury molecule-1 level are associated with adverse outcomes in acute renal failure. *J Am Soc Nephrol* 2007;18:904-912.
- Parr SK, Clark AJ, Bian A, et al. Urinary L-FABP predicts poor outcomes in critically ill patients with early acute kidney injury. *Kidney Int* 2015;87:640-648.