

21



**Fully Solved
Latest CBSE
Sample Paper**

issued on 16 September, 2022



**16 Fully
Solved Sample
Papers**



**5 Unsolved
Sample Papers with
Video Solution**

CBSE

Board Exams (2023)

MATHEMATICS (041) Class-12

O.P. GUPTA
INDIRA AWARD WINNER

PLEASURE TEST SERIES
SAMPLE QUESTION PAPERS

BASED ON THE LATEST CBSE XII
SYLLABUS & **LATEST PATTERN**
FOR CBSE XII BOARD EXAMS 2023



CBSE SAMPLE QUESTION PAPERS

(PLEASURE TEST SERIES)

CLASS XII MATHEMATICS 041

O.P. GUPTA

MATHS (H.), E & C ENGINEERING
INDIRA AWARD WINNER

© All Rights Reserved With Author (O.P. Gupta)
To Buy our **Best Seller** Math Books, WhatsApp @ +91 9650350480

Published by

THE O.P. GUPTA ADVANCED MATH CLASSES
1st Floor of 1625 D 4/A, Opp. HP Petrol Pump, Thana
Road, Najafgarh, New Delhi-43

Latest 2022-23 Edition Based on New Pattern

Friday, October 07, 2022

© All Rights Reserved With Author (O.P. GUPTA)
Copyright © Author

CAUTION

No part of this book or the complete book should be reproduced or copied in any form [photocopying, selling by any third party, resold, restored by information storage devices, or as the notes by any person(s)] without the prior consent of the author. Doing so will be considered as the intellectual theft and will deprive the author of his due credit for his work.

REMARK

While we've taken all possible care in the editing, proof reading and printing of this book, still some errors might have crept in. The author should not be held responsible for any misprint/omission. We shall feel grateful for the suggestions received from the readers for the further improvement of the book.

BOOKS BY O.P. GUPTA

MATHMISSION

For XII (2022-23)

MATHMISSION

For XI (2022-23)

MATHMISSION

Term II for XII (2021-22)

MATHMISSION

Term I for XII (2021-22)



O.P. GUPTA

Math Mentor

Indira Award Winner

MENTORS OF MATHMISSION

Mr Prabhat Marwaha

M.Sc. (Maths), B.Ed.

(32 Years Experience)

Vice Principal, Jawahar Navodaya Vidyalaya,
Pipersand, Lucknow

Mr Pankaj Chugh

M.Sc. (Maths), M.Phil. (Maths), B.Ed.

(31 Years Experience)

DAV Public School, Paschim Vihar
Edu-Maths, Sec. 18, Rohini, Delhi

Mr Amit Bajaj

M.Sc. (Maths), M.A. (Education), B.Ed.

(21 Years Experience)

Senior Maths Faculty
CRPF Public School, Rohini, Delhi

Ms Ashita Mehta

M.Sc. (Maths), B.Ed.

(16 Years Experience)


PGT (Maths)
GD Goenka Public School, Indirapuram, Ghaziabad

Mr Sachin Pandey

M.Sc. (Maths), B.Ed.

(14 Years Experience)

Coordinator
St. Mary's Sr. Sec. School, Rudrapur, Uttarakhand



A BRIEF SYNOPSIS Of CONTENTS IN

CBSE 21 SAMPLE PAPERS

For CBSE 2022-23 Exams ■ Class 12 Maths (041)

Pleasure Test Series By O.P. Gupta

- ✦ Multiple Choices Questions
- ✦ Assertion-Reason (A-R) Questions
- ✦ Subjective type Questions (2 Markers, 3 Markers & 5 Markers)
- ✦ CASE STUDY QUESTIONS (As per Latest format for 2023)
- ✦ H.O.T.S. Questions
- ✦ Detailed Solutions of 16 Sample Papers
- ✦ ANSWERS of 5 Unsolved Sample Papers

Most of the Pleasure Tests (PTS) are based on the Blueprint - same as that of CBSE Official Sample Question Paper. Though, in some of the PTS we have adopted different Blueprint : keeping in mind that the Unitwise weightage is not altered.

📺 For latest Math-Lectures, visit on
YouTube.com/MathematiciaByOPGupta

**For order related queries, please contact by
WhatsApp @ +91 9650350480 (only message)**

SHARE WITH OTHER STUDENTS ALSO TO HELP THEM IN THEIR PREPARATION.

TABLE OF CONTENTS

S. No.	TOPICS	PAGES
01.	Bifurcation of CBSE Sample Paper (2022-23)	
02.	CBSE SAMPLE PAPER (Issued on 16/09/2022)	(01 - 18)
03.	Sample Papers (with Detailed Solutions)	
	▪ PTS-01 to PTS-15	(19 - 285)
04.	Sample Papers (with Answers only)	(286 - 317)
	▪ PTS-16 to PTS-20	
05.	Reviews for the MATHMISSION Book	




Syllabus
CBSE EXAMS (2022-23)
Class XII ▪ Maths (041)

One Paper (Theory)
180 Minutes

Max Marks: 80

No.	UNITS	MARKS
I	Relations & Functions	08
II	Algebra	10
III	Calculus	35
IV	Vectors & 3 D Geometry	14
V	Linear Programming	05
VI	Probability	08
	Total	80

🔗 **OFFICIAL SAMPLE QUESTION PAPER & Unsolved Sample Papers (PTS-16 to PTS-20)** shall be Solved on YouTube Channel - **MATHEMATICIA By O.P. GUPTA**



OUR BOOKS HAVE GONE TO VARIOUS STATES OF INDIA & ABROAD

- Jammu & Kashmir
- Himachal Pradesh
- Punjab
- Chandigarh
- Rajasthan
- Delhi
- Haryana
- Uttarakhand
- Uttar Pradesh
- Bihar
- Jharkhand
- Odisha
- West Bengal

- Assam
- Tripura
- Madhya Pradesh
- Chhattisgarh
- Gujarat
- Telangana
- Andhra Pradesh
- Maharashtra
- Karnataka
- Tamilnadu
- Kerala
- Puducherry

OUR BOOKS HAVE ALSO BEEN TO FOREIGN COUNTRIES


- Oman
- Doha (Qatar)
- Saudi Arabia

MATHEMATICIA By O.P. GUPTA




Respected Teachers & dear Students

Over the years, I have been invited to address many sessions in schools, institutions for the **enrichment of students** across the Country and outside India as well. I've been invited as **Judge** also for Math events in various schools viz. Blue Bells Group of Schools etc. If you wish to invite me for such type of sessions, please contact us at below mentioned email.



Also if you feel that my work in Math has helped you in anyway, please **Share your feedback** in **written** or in **Video-mode**. It gives boost to work with more strength and determination for the Math community.



Let's spread a word about our work in Math - to help all the Students out there!

91-9650350480
iMathematicia@gmail.com

CBSE SQP • MATHEMATICS (041) • XII

By O.P. GUPTA, Indira Award Winner

Chapters	Section A (1 mark) <i>MCQ type</i>	Section B (2 marks) <i>VSA type</i>	Section C (3 marks) <i>SA type</i>	Section D (5 marks) <i>LA type</i>	Section E (Case Study) (4 marks) <i>Subjective type</i>	Marks for each unit
Relations & Functions		Q21*		Q33*		8
Inverse Trig. Functions	Q19 (AR)	Q21*				
Matrices & Determinants	Q01, 02, 10, 12, 13			Q35		10
Continuity & Differentiability	Q04, 16	Q24				
Applications of Derivatives		Q22			Q36*, Q37* (with 3 parts each)	35
Integrals	Q05, 09		Q26, Q28*, Q31			
Application of Integrals				Q32		
Differential Equations	Q06, 15		Q29*			
Vector Algebra	Q03, 08, 17	Q23*, Q25				14
3 Dimensional Geometry	Q18, 20 (AR)	Q23*		Q34*		
Linear Programming	Q07, 11		Q30			5
Probability	Q14		Q27*		Q38 (with 2 parts)	8
Total Marks	20 Marks	10 Marks	18 Marks	20 Marks	12 Marks	80 Marks

Note : This Bifurcation of Questions is based on Sample Question Paper issued by CBSE, for the Board Examination 2023.

** Internal choices given for the questions based on these topics / units.*

PLEASURE TEST SERIES PTS-19

A Compilation by
O.P. GUPTA, Math Mentor & Author
INDIRA AWARD WINNER

General Instructions : Same as given in PTS-01.

SECTION A

(Question numbers 01 to 20 carry 1 mark each.)

Followings are **multiple choice questions**. Select the correct option in each one of them.

01. If $A = [a_{ij}]_{m \times n}$ is a square matrix, then which of the following is true?

- (a) $m < n$ (b) $m > n$ (c) $m = n$ (d) $m = 0$

02. If $A = \begin{pmatrix} 1 & 0 & 0 \\ 0 & \cos \theta & \sin \theta \\ 0 & \sin \theta & -\cos \theta \end{pmatrix}$, then which of the following statements are correct?

I : $A + \text{adj.}A$ is a null matrix.

II : $A^{-1} + \text{adj.}A$ is a null matrix.

III : $A - A^{-1}$ is a null matrix.

Select the correct answer using the options given below.

- (a) I and II only (b) II and III only (c) I and III only (d) I, II and III all three

03. For $|\vec{a}| = 1$, $|\vec{b}| = 2$ and $\theta = \frac{\pi}{3}$ (where θ is angle between \vec{a} and \vec{b}), value of $\vec{a} \cdot \vec{b}$ is

- (a) 0 (b) -1 (c) 1 (d) 2

04. The value of $(-m)$ for which the function $f(x) = \begin{cases} \frac{1-e^{4x}}{x}, & \text{if } x \neq 0 \\ m, & \text{if } x = 0 \end{cases}$ is continuous at $x = 0$, is

- (a) -4 (b) 4 (c) 1 (d) 2

05. $\int (x^x)^2 (1 + \log x) dx =$

- (a) $x^{2x} + C$ (b) $\frac{x^{2x}}{2} + C$ (c) $2x^{2x} + C$ (d) $\frac{x^x}{2} + C$

06. Order of the differential equation $\left(\frac{d^2y}{dx^2}\right)^3 + \cos\left(\frac{dy}{dx}\right) = 0$ is

- (a) 1 (b) 2 (c) 3 (d) 4

07. For what value of x , the scalar projection of $\hat{i} - \hat{j} + x\hat{k}$ on $2\hat{i} + 2\hat{j} - 3\hat{k}$ is $\frac{6}{\sqrt{17}}$?

- (a) 2 (b) $\frac{1}{7}$ (c) 1 (d) -2

08. The value of $\int_0^1 \log\left(\frac{1}{x} - 1\right) dx$ is


- (a) 1 (b) 0 (c) -1 (d) $\log 2$

09. Let Δ be the determinant of a matrix A , where $A = \begin{pmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{pmatrix}$ and C_{11}, C_{12}, C_{13} be the

cofactors of a_{11}, a_{12}, a_{13} respectively. What is the value of $a_{11}C_{11} + a_{12}C_{12} + a_{13}C_{13}$?

- (a) 0 (b) 1 (c) Δ (d) $-\Delta$

10. For an L.P.P. the objective function is $Z = 4x + 3y$, and the feasible region determined by a set of constraints (linear inequations) is shown in the graph.



Which one of the following statements is true?

- (a) Maximum value of Z is at R
 (b) Maximum value of Z is at Q
 (c) Value of Z at R is less than the value at P
 (d) Value of Z at Q is less than the value of R

11. A linear programming problem is as follows :

Minimize : $z = 2x + y$

Subject to the constraints : $x \geq 3, x \leq 9, y \geq 0, x - y \geq 0, x + y \leq 14$.

The feasible region has

- (a) 5 corner points including $(0, 0)$ and $(9, 5)$
 (b) 5 corner points including $(7, 7)$ and $(3, 3)$
 (c) 5 corner points including $(14, 0)$ and $(9, 0)$
 (d) 5 corner points including $(3, 6)$ and $(9, 5)$

12. Area of a triangle is given by $\Delta = \frac{1}{2} \begin{vmatrix} 1 & 2 & 1 \\ 0 & 1 & 1 \\ 3 & k & 1 \end{vmatrix}$. If the area equals 1 Sq. unit then, $k =$

- (a) 0, 1 (b) 0, 2 (c) 2, 6 (d) 2, -6

13. Let $|A| = \begin{vmatrix} 1 & 0 & 0 \\ 1 & 2 & 3 \\ 0 & 4 & 8 \end{vmatrix}$. Then, $|\text{adj.} A| =$

- (a) 4 (b) 2 (c) 16 (d) ± 4

14. Given two independent events A and B such that $P(A) = 0.3, P(B) = 0.6$. Then $P(A | B)$ is

- (a) 0.9 (b) 0.18 (c) 0.28 (d) 0.3

15. To solve the D.E. $2y e^{x/y} dx + (y - 2x e^{x/y}) dy = 0$, we must substitute

- (a) $x = v$ (b) $x = vy$ (c) $e^x = v$ (d) $y = v$

16. If $y = e^{\tan^{-1} x}$, then $(1 + x^2)^2 y_2 + 2x(1 + x^2) y_1 =$

- (a) 0 (b) $-y$ (c) y (d) 2

17. Let \vec{a} and \vec{b} are two unit vectors such that $\vec{a} + 2\vec{b}$ and $5\vec{a} - 4\vec{b}$ are perpendicular. The angle θ between \vec{a} and \vec{b} is
- (a) $\frac{\pi}{6}$ (b) $\frac{\pi}{4}$ (c) $\frac{\pi}{3}$ (d) $\frac{\pi}{2}$
18. If a line makes angle $\frac{\pi}{2}$, $\frac{\pi}{3}$ and $\frac{\pi}{6}$ with the positive direction of x, y and z-axis respectively, then its direction cosines are
- (a) $0, \frac{\sqrt{3}}{2}, \frac{1}{2}$ (b) $0, \sqrt{3}, 1$ (c) $0, 1, \sqrt{3}$ (d) $0, \frac{1}{2}, \frac{\sqrt{3}}{2}$

Followings are **Assertion-Reason based questions**.

In the following questions, a statement of Assertion (A) is followed by a statement of Reason (R). Choose the correct answer out of the following choices.

- (a) Both A and R are true and R is the correct explanation of A.
 (b) Both A and R are true and R is not the correct explanation of A.
 (c) A is true but R is false.
 (d) A is false but R is true.
19. **Assertion (A) :** The value of $\tan^{-1}\{\cot(\operatorname{cosec}^{-1}2)\}$ is $\frac{\pi}{3}$.
- Reason (R) :** If $y = \sin^{-1}x$, then $y \in \left[-\frac{\pi}{2}, \frac{\pi}{2}\right]$.
20. **Assertion (A) :** The acute angle between the line $\vec{r} = \hat{i} + \hat{j} + 2\hat{k} + \lambda(\hat{i} - \hat{j})$ and the y-axis is $\frac{\pi}{2}$.
- Reason (R) :** The acute angle θ between the lines $\vec{r} = x_1\hat{i} + y_1\hat{j} + z_1\hat{k} + \lambda(a_1\hat{i} + b_1\hat{j} + c_1\hat{k})$ and $\vec{r} = x_2\hat{i} + y_2\hat{j} + z_2\hat{k} + \mu(a_2\hat{i} + b_2\hat{j} + c_2\hat{k})$ is given by $\cos\theta = \frac{|a_1a_2 + b_1b_2 + c_1c_2|}{\sqrt{a_1^2 + b_1^2 + c_1^2}\sqrt{a_2^2 + b_2^2 + c_2^2}}$.

SECTION B

(Question numbers 21 to 25 carry 2 marks each.)

21. Find the value of $\cos^{-1}\left[\cos\left(\frac{13\pi}{8}\right)\right]$.

OR

Let $A = \{m, a, t, h\}$. Write the number of reflexive relations defined on A.

Also, write the number of symmetric relations defined on A.

22. The volume of a cone is changing at the rate of $40 \text{ cm}^3/\text{s}$. If height of the cone is always equal to its diameter, then find the rate of change of radius when its circular base area is 1 square metre.
23. The position vectors of vertices A, B and C of triangle ABC are respectively $\hat{j} + \hat{k}$, $3\hat{i} + \hat{j} + 5\hat{k}$ and $3\hat{j} + 3\hat{k}$. Find $\angle BCA$ of triangle ABC.

OR

Using the concept of direction ratios of line, show that the points A(2, 3, -4), B(1, -2, 3) and C(3, 8, -11) are collinear.

24. If $\sin y = x \sin(a + y)$ then, find $\sin a \frac{d^2y}{dx^2} - \sin 2(a + y) \frac{dy}{dx}$.

25. If $\vec{a} = 2\hat{i} + \hat{j} - 2\hat{k}$ and $\vec{b} = 3\hat{i} - 5\hat{j} + 4\hat{k}$ then, find $\frac{\text{Vector projection of } \vec{a} \text{ on } \vec{b}}{\text{Projection of } \vec{b} \text{ on } \vec{a}}$.

SECTION C

(Question numbers 26 to 31 carry 3 marks each.)

26. Find : $\int \frac{(\cos x)^{3/2} - (\sin x)^{3/2}}{\sqrt{\sin x \cos x}} dx$.
27. A and B throw a die alternatively till one of them gets a 'six' and wins the game. Find their respective probabilities of winning, if A starts the game.

OR

There is a group of 50 people who are patriotic out of which 20 believe in non-violence. Two persons are selected at random out of them. Find the mean of the probability distribution for the selected persons who are non-violent.

28. Evaluate : $\int_0^{\pi} |\cos x - \sin x| dx$.

OR

Evaluate : $\int_0^{\pi/4} \log(1 + \tan x) dx$.

29. Solve the differential equation : $(dy - dx) + \cos x (dy + dx) = 0$.

OR

Solve the differential equation : $x \cos\left(\frac{y}{x}\right) \times \frac{dy}{dx} = y \cos\left(\frac{y}{x}\right) + x, x > 0$.

30. Solve the following Linear Programming Problem graphically :
Minimize $Z = 10x + 8y$ subject to $3x + y \geq 300, x + y \geq 240; x, y \geq 0$.

31. Find : $\int \sqrt{\frac{1-x}{1+x}} dx$.

SECTION D

(Question numbers 32 to 35 carry 5 marks each.)

32. Make a rough sketch of the region $\{(x, y) : 0 \leq y \leq x^2, 0 \leq y \leq x, 0 \leq x \leq 3\}$ and find the area of the region using integration.
33. Let $Y = \{n^2 : n \in \mathbb{N}\} \subset \mathbb{N}$. Consider $f : \mathbb{N} \rightarrow Y$ given as $f(n) = n^2$.
Show that the function f is one-one and onto.

OR

Show that the relation R in the set A of points in a plane given by $R = \{(P, Q) : \text{Distance of the point } P \text{ from the origin is same as the distance of point } Q \text{ from the origin}\}$, is an equivalence relation.

34. Find the shortest distance between the lines $\vec{r} = \hat{i} + \hat{j} + \lambda(2\hat{i} - \hat{j} + \hat{k})$ and $\vec{r} = 2\hat{i} + \hat{j} - \hat{k} + \mu(3\hat{i} - 5\hat{j} + 2\hat{k})$.
Hence, obtain the acute angle between the lines.

OR

A line passing through the point A with position vector $\vec{a} = 4\hat{i} + 2\hat{j} + 2\hat{k}$ is parallel to the vector $\vec{b} = 2\hat{i} + 3\hat{j} + 6\hat{k}$. Find the length of the perpendicular drawn on this line from a point P with the position vector $\vec{r}_1 = \hat{i} + 2\hat{j} + 3\hat{k}$.

35. Solve the system of equations $x + y + z = 21$, $4x + 3y + 2z = 60$, $6x + 2y + 3z = 70$, using matrices.

SECTION E

(Question numbers 36 to 38 carry 4 marks each.)

This section contains **three Case-study / Passage based questions**.

First two questions have **three sub-parts** (i), (ii) and (iii) of **marks 1, 1 and 2** respectively.

Third question has **two sub-parts** of **2 marks** each.

36. **CASE STUDY I :** Read the following passage and the answer the questions given below.



A shopkeeper sells three types of flowers seeds A_1 , A_2 and A_3 .

These are sold as mixture, where their proportions are 4:4:2 respectively.

Also their germination rates are 45%, 60% and 35% respectively.

Let A_1 : seed A_1 is chosen, A_2 : seed A_2 is chosen and A_3 : seed A_3 is chosen 3.

Also let E : seed germinates.

(i) Find $P(A_1)$, $P(A_2)$ and $P(A_3)$.


(ii) Write $P(E | A_1) + P(E | A_2) + P(E | A_3)$.

(iii) Calculate the probability of a randomly chosen seed to germinate. Express the answer in %.

OR

(iii) Calculate the probability that it is of the type A_2 given that a randomly chosen seed does not germinate.

37. **CASE STUDY II :** Read the following passage and answer the questions given below.



In an elliptical sport field the authority wants to design a rectangular soccer field with the


maximum possible area. The sport field is given by the graph of $\frac{x^2}{25} + \frac{y^2}{9} = 1$.

- (i) If the length and the breadth of the rectangular field be ' $2x$ ' and ' $2y$ ' respectively, then find the area function in terms of ' x '.
- (ii) Find the critical point of the function obtained in (i).
- (iii) Use first derivative test to find the length ' $2x$ ' and width ' $2y$ ' of the soccer field, that will maximize its area.

OR

- (iii) Use second derivative test to find the length ' $2x$ ' and width ' $2y$ ' of the soccer field, that will maximize its area.

38. **CASE STUDY III** : Read the following passage and answer the questions given below.



A cylindrical tank of radius 10 m is being filled with water at the rate of 314 cubic metre per hour.

- (i) Find the rate of increase of the depth (h) of the water in the tank. Use $\pi = 3.14$.
- (ii) Find the rate of increase of the curved surface area (A) of the water in the tank.

SAMPLE PAPER

issued by CBSE Exams (2022-23)
Mathematics (041) - Class 12

Time Allowed : 180 Minutes

Max. Marks : 80

General Instructions :

1. This Question paper contains **five sections - A, B, C, D and E**. Each section is compulsory. However, there are **internal choices** in some questions.
2. Section A has **18 MCQs** and **02 Assertion-Reason (A-R)** based questions of **1 mark** each.
Section B has **05 questions** of **2 marks** each.
Section C has **06 questions** of **3 marks** each.
Section D has **04 questions** of **5 marks** each.
Section E has **03 Case-study / Source-based / Passage-based** questions with **sub-parts (4 marks each)**.
3. There is no overall choice. However, **internal choice** has been provided in
 - **02 Questions of Section B**
 - **03 Questions of Section C**
 - **02 Questions of Section D**
 - **02 Questions of Section E**You have to attempt only one of the alternatives in all such questions.

SECTION A

(Question numbers 01 to 20 carry **1 mark** each.)


Followings are **multiple choice questions**. Select the correct option in each one of them.

01. If $A = [a_{ij}]$ is a skew-symmetric matrix of order n , then
 - (a) $a_{ij} = \frac{1}{a_{ji}} \forall i, j$
 - (b) $a_{ij} \neq 0 \forall i, j$
 - (c) $a_{ij} = 0$, where $i = j$
 - (d) $a_{ij} \neq 0$ where $i = j$
02. If A is a square matrix of order 3, $|A'| = -3$, then $|AA'| =$
 - (a) 9
 - (b) -9
 - (c) 3
 - (d) -3
03. The area of a triangle with vertices A, B, C is given by
 - (a) $|\overrightarrow{AB} \times \overrightarrow{AC}|$
 - (b) $\frac{1}{2} |\overrightarrow{AB} \times \overrightarrow{AC}|$
 - (c) $\frac{1}{4} |\overrightarrow{AC} \times \overrightarrow{AB}|$
 - (d) $\frac{1}{8} |\overrightarrow{AC} \times \overrightarrow{AB}|$
04. The value of 'k' for which the function $f(x) = \begin{cases} \frac{1 - \cos 4x}{8x^2}, & \text{if } x \neq 0 \\ k, & \text{if } x = 0 \end{cases}$ is continuous at $x = 0$, is
 - (a) 0
 - (b) -1
 - (c) 1
 - (d) 2
05. If $f'(x) = x + \frac{1}{x}$, then $f(x)$ is
 - (a) $x^2 + \log|x| + C$
 - (b) $\frac{x^2}{2} + \log|x| + C$
 - (c) $\frac{x}{2} + \log|x| + C$
 - (d) $\frac{x}{2} - \log|x| + C$

06. If m and n respectively, are the order and the degree of the differential equation

$$\frac{d}{dx} \left[\left(\frac{dy}{dx} \right)^4 \right] = 0, \text{ then } m + n =$$

- (a) 1 (b) 2 (c) 3 (d) 4
07. The solution set of the inequality $3x + 5y < 4$ is
 (a) an open half-plane not containing the origin
 (b) an open half-plane containing the origin
 (c) the whole XY -plane not containing the line $3x + 5y = 4$
 (d) a closed half-plane containing the origin
08. The scalar projection of the vector $3\hat{i} - \hat{j} - 2\hat{k}$ on the vector $\hat{i} + 2\hat{j} - 3\hat{k}$ is
 (a) $\frac{7}{\sqrt{14}}$ (b) $\frac{7}{14}$ (c) $\frac{6}{13}$ (d) $\frac{7}{2}$
09. The value of $\int_2^3 \frac{x}{x^2 + 1} dx$ is
 (a) $\log 4$ (b) $\log \frac{3}{2}$ (c) $\frac{1}{2} \log 2$ (d) $\log \frac{9}{4}$
10. If A, B are non-singular square matrices of the same order, then $(AB^{-1})^{-1} =$
 (a) $A^{-1}B$ (b) $A^{-1}B^{-1}$ (c) BA^{-1} (d) AB
11. The corner points of the shaded unbounded feasible region of an LPP are $(0, 4)$, $(0.6, 1.6)$ and $(3, 0)$ as shown in the figure.
 The minimum value of the objective function $Z = 4x + 6y$ occurs at



- (a) $(0.6, 1.6)$ only
 (b) $(3, 0)$ only
 (c) $(0.6, 1.6)$ and $(3, 0)$ only
 (d) every point of the line-segment joining the points $(0.6, 1.6)$ and $(3, 0)$
12. If $\begin{vmatrix} 2 & 4 \\ 5 & 1 \end{vmatrix} = \begin{vmatrix} 2x & 4 \\ 6 & x \end{vmatrix}$, then the possible value(s) of 'x' is/are
 (a) 3 (b) $\sqrt{3}$ (c) $-\sqrt{3}$ (d) $\sqrt{3}, -\sqrt{3}$
13. If A is a square matrix of order 3 and $|A| = 5$, then $|\text{adj } A| =$

- (a) 5 (b) 25 (c) 125 (d) $\frac{1}{5}$
14. Given two independent events A and B such that $P(A) = 0.3$, $P(B) = 0.6$ and $P(A' \cap B')$ is
(a) 0.9 (b) 0.18 (c) 0.28 (d) 0.1
15. The general solution of the differential equation $ydx - xdy = 0$ is
(a) $xy = C$ (b) $x = Cy^2$ (c) $y = Cx$ (d) $y = Cx^2$
16. If $y = \sin^{-1} x$, then $(1 - x^2)y_2$ is equal to
(a) xy_1 (b) xy (c) xy_2 (d) x^2
17. If two vectors \vec{a} and \vec{b} are such that $|\vec{a}| = 2$, $|\vec{b}| = 3$ and $\vec{a} \cdot \vec{b} = 4$, then $|\vec{a} - 2\vec{b}|$ is equal to
(a) $\sqrt{2}$ (b) $2\sqrt{6}$ (c) 24 (d) $2\sqrt{2}$
18. P is a point on the line joining the points A(0, 5, -2) and B(3, -1, 2). If the x-coordinate of P is 6, then its z-coordinate is
(a) 10 (b) 6 (c) -6 (d) -10

Followings are **Assertion-Reason based questions**.

In the following questions, a statement of Assertion (A) is followed by a statement of Reason (R).

Choose the correct answer out of the following choices.

- (a) Both A and R are true and R is the correct explanation of A.
(b) Both A and R are true and R is not the correct explanation of A.
(c) A is true but R is false.
(d) A is false but R is true.
19. **Assertion (A) :** The domain of the function $\sec^{-1} 2x$ is $\left(-\infty, -\frac{1}{2}\right] \cup \left[\frac{1}{2}, \infty\right)$.
Reason (R) : $\sec^{-1}(-2) = -\frac{\pi}{4}$.
20. **Assertion (A) :** The acute angle between the line $\vec{r} = \hat{i} + \hat{j} + 2\hat{k} + \lambda(\hat{i} - \hat{j})$ and the x-axis is $\frac{\pi}{4}$.
Reason (R) : The acute angle θ between the lines
 $\vec{r} = x_1\hat{i} + y_1\hat{j} + z_1\hat{k} + \lambda(a_1\hat{i} + b_1\hat{j} + c_1\hat{k})$ and
 $\vec{r} = x_2\hat{i} + y_2\hat{j} + z_2\hat{k} + \mu(a_2\hat{i} + b_2\hat{j} + c_2\hat{k})$ is given by $\cos \theta = \frac{|a_1a_2 + b_1b_2 + c_1c_2|}{\sqrt{a_1^2 + b_1^2 + c_1^2} \sqrt{a_2^2 + b_2^2 + c_2^2}}$.

SECTION B

(Question numbers 21 to 25 carry 2 marks each.)

21. Find the value of $\sin^{-1} \left[\sin \left(\frac{13\pi}{7} \right) \right]$.

OR

Prove that the function f is surjective, where $f : \mathbb{N} \rightarrow \mathbb{N}$ such that

$$f(n) = \begin{cases} \frac{n+1}{2}, & \text{if } n \text{ is odd} \\ \frac{n}{2}, & \text{if } n \text{ is even} \end{cases}$$

Is the function injective? Justify your answer.

22. A man 1.6 m tall walks at the rate of 0.3 m/sec away from a street light that is 4 m above the ground. At what rate is the tip of his shadow moving? At what rate is his shadow lengthening?
23. If $\vec{a} = \hat{i} - \hat{j} + 7\hat{k}$ and $\vec{b} = 5\hat{i} - \hat{j} + \lambda\hat{k}$, then find the value of λ so that the vectors $\vec{a} + \vec{b}$ and $\vec{a} - \vec{b}$ are orthogonal.

OR

Find the direction ratios and direction cosines of a line parallel to the line whose equation is $6x - 12 = 3y + 9 = 2z - 2$.

24. If $y\sqrt{1-x^2} + x\sqrt{1-y^2} = 1$, then prove that $\frac{dy}{dx} = -\sqrt{\frac{1-y^2}{1-x^2}}$.
25. Find $|\vec{x}|$, if $(\vec{x} - \vec{a}) \cdot (\vec{x} + \vec{a}) = 12$, where \vec{a} is a unit vector.

SECTION C

(Question numbers 26 to 31 carry 3 marks each.)

26. Find : $\int \frac{dx}{\sqrt{3-2x-x^2}}$.
27. Three friends go for coffee. They decide who will pay the bill, by each tossing a coin and then letting the "odd person" pay. There is no odd person if all three tosses produce the same result. If there is no odd person in the first round, they make a second round of tosses and they continue to do so until there is an odd person. What is the probability that exactly three rounds of tosses are made?

OR

Find the mean number of defective items in a sample of two items drawn one-by-one without replacement from an urn containing 6 items, which include 2 defective items. Assume that the items are identical in shape and size.

28. Evaluate : $\int_{\pi/6}^{\pi/3} \frac{dx}{1 + \sqrt{\tan x}}$.

OR

Evaluate : $\int_0^4 |x-1| dx$.

29. Solve the differential equation : $ydx + (x - y^2)dy = 0$.

OR

Solve the differential equation : $xdy - ydx = \sqrt{x^2 + y^2} dx$.

30. Solve the following Linear Programming Problem graphically :
Maximize $Z = 400x + 300y$ subject to $x + y \leq 200$, $x \leq 40$, $x \geq 20$, $y \geq 0$.
31. Find : $\int \frac{(x^3 + x + 1)}{(x^2 - 1)} dx$.

SECTION D

(Question numbers 32 to 35 carry 5 marks each.)

32. Make a rough sketch of the region $\{(x, y) : 0 \leq y \leq x^2, 0 \leq y \leq x, 0 \leq x \leq 2\}$ and find the area of the region using integration.
33. Define the relation R in the set $N \times N$ as follows :
For $(a, b), (c, d) \in N \times N$, $(a, b) R (c, d)$ iff $ad = bc$.
Prove that R is an equivalence relation in $N \times N$.

OR

Given a non-empty set X , define the relation R in $P(X)$ as follows :

For $A, B \in P(X)$, $(A, B) \in R$ iff $A \subset B$.

Prove that R is reflexive, transitive and not symmetric.

34. An insect is crawling along the line $\vec{r} = 6\hat{i} + 2\hat{j} + 2\hat{k} + \lambda(\hat{i} - 2\hat{j} + 2\hat{k})$ and another insect is crawling along the line $\vec{r} = -4\hat{i} - \hat{k} + \mu(3\hat{i} - 2\hat{j} - 2\hat{k})$. At what points on the lines should they reach so that the distance between them is the shortest? Find the shortest possible distance between them.

OR

The equations of motion of a rocket are :

$x = 2t$, $y = -4t$, $z = 4t$, where the time t is given in seconds, and the coordinates of a moving point in km. What is the path of the rocket? At what distances will the rocket be from the starting point $O(0, 0, 0)$ and from the following line in 10 seconds?

$$\vec{r} = 20\hat{i} - 10\hat{j} + 40\hat{k} + \mu(10\hat{i} - 20\hat{j} + 10\hat{k}).$$

35. If $A = \begin{bmatrix} 2 & -3 & 5 \\ 3 & 2 & -4 \\ 1 & 1 & -2 \end{bmatrix}$, find A^{-1} . Use A^{-1} to solve the following system of equations

$$2x - 3y + 5z = 11, 3x + 2y - 4z = -5, x + y - 2z = -3.$$

SECTION E

(Question numbers 36 to 38 carry 4 marks each.)

This section contains **three Case-study / Passage based questions**.

First two questions have **three sub-parts** (i), (ii) and (iii) of **marks 1, 1 and 2** respectively.

Third question has **two sub-parts** of **2 marks** each.

36. **CASE STUDY I :** Read the following passage and the answer the questions given below.



The temperature of a person during an intestinal illness is given by

$f(x) = -0.1x^2 + mx + 98.6$, $0 \leq x \leq 12$, m being a constant, where $f(x)$ is the temperature in $^{\circ}\text{F}$ at x days.

- Is the function differentiable in the interval $(0, 12)$? Justify your answer.
- If 6 is the critical point of the function, then find the value of the constant m .
- Find the intervals in which the function is strictly increasing / strictly decreasing.

OR

- Find the points of local maximum / local minimum, if any, in the interval $(0, 12)$ as well as the points of absolute maximum / absolute minimum in the interval $[0, 12]$. Also, find the

corresponding local maximum / local minimum and the absolute maximum / absolute minimum values of the function.

37. **CASE STUDY II :** Read the following passage and answer the questions given below.



In an elliptical sport field the authority wants to design a rectangular soccer field with the maximum possible area. The sport field is given by the graph of $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$.

- (i) If the length and the breadth of the rectangular field be $2x$ and $2y$ respectively, then find the area function in terms of x .
- (ii) Find the critical point of the function.
- (iii) Use First Derivative test to find the length $2x$ and width $2y$ of the soccer field (in terms of a and b) that maximize its area.

OR

- (iii) Use Second Derivative test to find the length $2x$ and width $2y$ of the soccer field (in terms of a and b) that maximize its area.

38. **CASE STUDY III :** Read the following passage and answer the questions given below.



There are two anti craft guns, named as A and B. The probabilities that the shell fired from them hits an airplane are 0.3 and 0.2 respectively. Both of them fired one shell at an airplane at the same time.

- (i) What is the probability that the shell fired from exactly one of them hit the plane?
- (ii) If it is known that the shell fired from exactly one of them hit the plane, then what is the probability that it was fired from B?



SOLUTIONS Of CBSE SQP-2023

SECTION A

For Class XII • Mathematics (041)

01. (c) In a skew-symmetric matrix, $a_{ij} = -a_{ji}$. Hence for $i = j$, we have $a_{ii} = 0$.

02. (a) $|AA'| = |A||A'| = |A||A| = (-3)(-3) = 9$.

03. (b) The area of the parallelogram with adjacent sides \overrightarrow{AB} and $\overrightarrow{AC} = |\overrightarrow{AB} \times \overrightarrow{AC}|$.

Hence, the area of the triangle with vertices $A, B, C = \frac{1}{2} |\overrightarrow{AB} \times \overrightarrow{AC}|$.

04. (c) The function f is continuous at $x = 0$, if $\lim_{x \rightarrow 0} f(x) = f(0) \dots (i)$

We have $f(0) = k$ and $\lim_{x \rightarrow 0} f(x) = \lim_{x \rightarrow 0} \frac{1 - \cos 4x}{8x^2} = \lim_{x \rightarrow 0} \frac{2\sin^2 2x}{8x^2} = \lim_{x \rightarrow 0} \frac{\sin^2 2x}{4x^2}$

$$\Rightarrow \lim_{x \rightarrow 0} \left(\frac{\sin 2x}{2x} \right)^2 = 1$$

Hence by (i), $k = 1$.

05. (b) $\therefore f(x) = \int f'(x) dx \quad \therefore f(x) = \int \left(x + \frac{1}{x} \right) dx$

So, $f(x) = \frac{x^2}{2} + \log|x| + C$.

06. (c) The given differential equation is $4 \left(\frac{dy}{dx} \right)^3 \left(\frac{d^2y}{dx^2} \right) = 0$. Here, $m = 2$ and $n = 1$.

Hence, $m + n = 3$.

07. (b) The strict inequality represents an open half plane and it contains the origin as $(0, 0)$ satisfies the inequality $3x + 5y < 4$.

08. (a) Scalar Projection of $3\hat{i} - \hat{j} - 2\hat{k}$ on vector $\hat{i} + 2\hat{j} - 3\hat{k} = \frac{(3\hat{i} - \hat{j} - 2\hat{k}) \cdot (\hat{i} + 2\hat{j} - 3\hat{k})}{|\hat{i} + 2\hat{j} - 3\hat{k}|} = \frac{7}{\sqrt{14}}$.

09. (c) $\int_2^3 \frac{x}{x^2 + 1} dx = \frac{1}{2} \int_2^3 \frac{2x}{x^2 + 1} dx = \frac{1}{2} [\log(x^2 + 1)]_2^3 = \frac{1}{2} (\log 10 - \log 5) = \frac{1}{2} \log \left(\frac{10}{5} \right) = \frac{1}{2} \log 2$.

10. (c) $(AB^{-1})^{-1} = (B^{-1})^{-1} A^{-1} = BA^{-1}$.

11. (d) The minimum value of the objective function occurs at two adjacent corner points $(0.6, 1.6)$ and $(3, 0)$ and there is no point in the half plane $4x + 6y < 12$ which is common with the feasible region. So, the minimum value occurs at every point of the line segment joining the two points.

12. (d) Note that, $2 \times 1 - 5 \times 4 = 2x^2 - 24$

$$\Rightarrow 2x^2 = 6 \quad \Rightarrow x^2 = 3$$

$$\Rightarrow x = \pm\sqrt{3}$$

13. (b) $\therefore |\text{adj } A| = |A|^{n-1} \quad \therefore |\text{adj } A| = |A|^{3-1}$

$$\Rightarrow |\text{adj } A| = 25$$

14. (c) $P(A' \cap B') = P(A') \times P(B')$ (As A and B are independent, A' and B' are also independent.)
 $= 0.7 \times 0.4 = 0.28$.

15. (c) $ydx - xdy = 0 \quad \Rightarrow ydx = xdy$

$$\Rightarrow \int \frac{dy}{y} = \int \frac{dx}{x}$$

$$\Rightarrow \log|y| = \log|x| + \log K, K > 0$$

$$\Rightarrow \log|y| = \log[|x|K]$$

$$\Rightarrow |y| = |x|K \Rightarrow y = \pm Kx$$

$$\therefore y = Cx, \text{ where } C = \pm K.$$

16. (a) $y = \sin^{-1} x \Rightarrow \frac{dy}{dx} = \frac{1}{\sqrt{1-x^2}}$

$$\Rightarrow \sqrt{1-x^2} \cdot \frac{dy}{dx} = 1$$

Again, differentiating both sides w.r. to x , we get $\sqrt{1-x^2} \frac{d^2y}{dx^2} + \frac{dy}{dx} \cdot \left(\frac{-2x}{2\sqrt{1-x^2}} \right) = 0$

On simplifying, we get $(1-x^2)y_2 = xy_1$.

17. (b) $|\vec{a} - 2\vec{b}|^2 = (\vec{a} - 2\vec{b}) \cdot (\vec{a} - 2\vec{b})$

$$\Rightarrow |\vec{a} - 2\vec{b}|^2 = \vec{a} \cdot \vec{a} - 4\vec{a} \cdot \vec{b} + 4\vec{b} \cdot \vec{b}$$

$$\Rightarrow |\vec{a} - 2\vec{b}|^2 = |\vec{a}|^2 - 4\vec{a} \cdot \vec{b} + 4|\vec{b}|^2$$

$$\Rightarrow |\vec{a} - 2\vec{b}|^2 = 4 - 16 + 36 = 24$$

$$\therefore |\vec{a} - 2\vec{b}| = 2\sqrt{6}.$$

18. (b) The line through the points $(0, 5, -2)$ and $(3, -1, 2)$ is $\frac{x}{3-0} = \frac{y-5}{-1-5} = \frac{z+2}{2+2}$

i.e., $\frac{x}{3} = \frac{y-5}{-6} = \frac{z+2}{4} = k$ (say)

Any point on the line is $P(3k, -6k+5, 4k-2)$, where k is an arbitrary scalar.

As x -coordinate of point P is 6 so, $3k = 6 \therefore k = 2$.

The z -coordinate of the point P will be $4 \times 2 - 2 = 6$.

19. (c) Recall that, $\sec^{-1} x$ is defined if $x \leq -1$ or $x \geq 1$.

Hence, $\sec^{-1} 2x$ will be defined if $x \leq -\frac{1}{2}$ or $x \geq \frac{1}{2}$.

So, A is true.

Also note that, $\sec^{-1}(-2) = \frac{2\pi}{3}$.

So, R is false.

20. (a) The equation of the x -axis may be written as $\vec{r} = \mu(\hat{i})$. Hence, the acute angle θ between the given line and the x -axis is given by $\cos \theta = \frac{|1 \times 1 + (-1) \times 0 + 0 \times 0|}{\sqrt{1^2 + (-1)^2 + 0^2} \sqrt{1^2 + 0^2 + 0^2}} = \frac{1}{\sqrt{2}} \Rightarrow \theta = \frac{\pi}{4}$.

So, A is true. Similarly R is true. Also R is the correct explanation of A.

SECTION B

21. $\sin^{-1} \left[\sin \left(\frac{13\pi}{7} \right) \right] = \sin^{-1} \left[\sin \left(2\pi - \frac{\pi}{7} \right) \right] = \sin^{-1} \left[-\sin \left(\frac{\pi}{7} \right) \right] = \sin^{-1} \left[\sin \left(-\frac{\pi}{7} \right) \right] = -\frac{\pi}{7}$.

OR

Let $y \in \mathbb{N}$ (codomain). Then $\exists 2y \in \mathbb{N}$ (domain) such that

$$f(2y) = \frac{2y}{2} = y. \text{ Hence, } f \text{ is surjective.}$$

Note that $1, 2 \in \mathbb{N}$ (domain) such that $f(1) = 1 = f(2)$.


Hence, f is not injective.

22. Let AB represent the height of the street light from the ground. At any time t seconds, let the man represented as ED of height 1.6 m be at a distance of x m from AB and the length of his shadow EC by y m.

Using similarity of triangles, we've $\frac{4}{1.6} = \frac{x+y}{y} \Rightarrow 3y = 2x$

Differentiating both sides w.r. to t , we get $3 \times \frac{dy}{dt} = 2 \times \frac{dx}{dt}$

$$\Rightarrow \frac{dy}{dt} = \frac{2}{3} \times 0.3 = 0.2.$$



At any time t seconds, the tip of his shadow is at a distance of $(x+y)$ m from AB .

The rate at which the tip of his shadow moving = $\left(\frac{dx}{dt} + \frac{dy}{dt} \right) \text{ m/s} = (0.3 + 0.2) \text{ m/s} = 0.5 \text{ m/s}.$

Also, the rate at which his shadow is lengthening = $\frac{dy}{dt} \text{ m/s} = 0.2 \text{ m/s}.$

23. $\vec{a} = \hat{i} - \hat{j} + 7\hat{k}$ and $\vec{b} = 5\hat{i} - \hat{j} + \lambda\hat{k}$

Hence, $\vec{a} + \vec{b} = 6\hat{i} - 2\hat{j} + (7+\lambda)\hat{k}$ and $\vec{a} - \vec{b} = -4\hat{i} + (7-\lambda)\hat{k}$

Now $\vec{a} + \vec{b}$ and $\vec{a} - \vec{b}$ will be orthogonal if, $(\vec{a} + \vec{b}) \cdot (\vec{a} - \vec{b}) = 0$

i.e., if, $-24 + (49 - \lambda^2) = 0 \Rightarrow \lambda^2 = 25$

i.e., if, $\lambda = \pm 5.$

OR

Equation of the line $6x - 12 = 3y + 9 = 2z - 2$, which when written in standard symmetric form,

will be $\frac{x-2}{\frac{1}{6}} = \frac{y-(-3)}{\frac{1}{3}} = \frac{z-1}{\frac{1}{2}}$

Since, lines are parallel, we have $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$

Hence, the required direction ratios are $\frac{1}{6}, \frac{1}{3}, \frac{1}{2}$ i.e., 1, 2, 3.

And the required direction cosines are $\frac{1}{\sqrt{14}}, \frac{2}{\sqrt{14}}, \frac{3}{\sqrt{14}}.$

24. Let $\sin^{-1} x = A$ and $\sin^{-1} y = B$. Then $x = \sin A$ and $y = \sin B$

So, $y\sqrt{1-x^2} + x\sqrt{1-y^2} = 1$ implies, $\sin B \cos A + \sin A \cos B = 1$

$\Rightarrow \sin(A+B) = 1$

$\Rightarrow A+B = \sin^{-1} 1 = \frac{\pi}{2}$

$\Rightarrow \sin^{-1} x + \sin^{-1} y = \frac{\pi}{2}$

Differentiating w.r. to x , we obtain $\frac{1}{\sqrt{1-x^2}} + \frac{1}{\sqrt{1-y^2}} \times \frac{dy}{dx} = 0$

$$\Rightarrow \frac{dy}{dx} = -\sqrt{\frac{1-y^2}{1-x^2}}.$$

25. Since \vec{a} is a unit vector, $\therefore |\vec{a}| = 1$

$$\text{As } (\vec{x} - \vec{a}) \cdot (\vec{x} + \vec{a}) = 12$$

$$\Rightarrow \vec{x} \cdot \vec{x} + \vec{x} \cdot \vec{a} - \vec{a} \cdot \vec{x} - \vec{a} \cdot \vec{a} = 12$$

$$\Rightarrow |\vec{x}|^2 - |\vec{a}|^2 = 12.$$

$$\Rightarrow |\vec{x}|^2 - 1^2 = 12$$

$$\Rightarrow |\vec{x}|^2 = 13$$

$$\therefore |\vec{x}| = \sqrt{13}.$$

SECTION C

26.
$$\int \frac{dx}{\sqrt{3-2x-x^2}}$$

$$= \int \frac{dx}{\sqrt{-(x^2+2x-3)}} = \int \frac{dx}{\sqrt{4-(x+1)^2}} = \int \frac{dx}{\sqrt{2^2-(x+1)^2}}$$

$$= \sin^{-1}\left(\frac{x+1}{2}\right) + C.$$

27. P (not obtaining an odd person in a single round)
 = P (all three of them throw tails or all three of them throw heads)
 $= \frac{1}{2} \times \frac{1}{2} \times \frac{1}{2} \times 2 = \frac{1}{4}$

P (obtaining an odd person in a single round)

$$= 1 - P(\text{not obtaining an odd person in a single round}) = \frac{3}{4}.$$

The required probability

$$= P\left(\begin{array}{l} \text{'in first round there is no odd person' and 'in second round there} \\ \text{is no odd person' and 'in third round there is an odd person'} \end{array}\right)$$

$$= \frac{1}{4} \times \frac{1}{4} \times \frac{3}{4} = \frac{3}{64}.$$

OR

Let X denote the Random Variable defined by the number of defective items.

$$P(X=0) = \frac{4}{6} \times \frac{3}{5} = \frac{2}{5}, P(X=1) = 2 \times \left(\frac{2}{6} \times \frac{4}{5}\right) = \frac{8}{15}, P(X=2) = \frac{2}{6} \times \frac{1}{5} = \frac{1}{15}.$$

The table for probability distribution is given below.

x_i	0	1	2
p_i	$\frac{6}{15}$	$\frac{8}{15}$	$\frac{1}{15}$

$$\text{Mean} = \sum p_i x_i = 0 + \frac{8}{15} + \frac{2}{15} = \frac{10}{15} = \frac{2}{3}.$$

28. Let $I = \int_{\pi/6}^{\pi/3} \frac{dx}{1 + \sqrt{\tan x}} = \int_{\pi/6}^{\pi/3} \frac{\sqrt{\cos x}}{\sqrt{\sin x} + \sqrt{\cos x}} dx \dots (i)$

Using $\int_a^b f(x) dx = \int_a^b f(a + b - x) dx$, we get

$$I = \int_{\pi/6}^{\pi/3} \frac{\sqrt{\cos\left(\frac{\pi}{6} + \frac{\pi}{3} - x\right)}}{\sqrt{\sin\left(\frac{\pi}{6} + \frac{\pi}{3} - x\right)} + \sqrt{\cos\left(\frac{\pi}{6} + \frac{\pi}{3} - x\right)}} dx$$

$$\Rightarrow I = \int_{\pi/6}^{\pi/3} \frac{\sqrt{\sin x}}{\sqrt{\cos x} + \sqrt{\sin x}} dx \dots (ii)$$

Adding (i) and (ii), we get

$$2I = \int_{\pi/6}^{\pi/3} \frac{\sqrt{\cos x}}{\sqrt{\sin x} + \sqrt{\cos x}} dx + \int_{\pi/6}^{\pi/3} \frac{\sqrt{\sin x}}{\sqrt{\cos x} + \sqrt{\sin x}} dx$$

$$\Rightarrow 2I = \int_{\pi/6}^{\pi/3} dx$$

$$\Rightarrow 2I = [x]_{\pi/6}^{\pi/3} = \frac{\pi}{3} - \frac{\pi}{6} = \frac{\pi}{6}$$

Hence, $I = \frac{\pi}{12}$.

OR

$$\begin{aligned} \int_0^4 |x-1| dx &= \int_0^1 |x-1| dx + \int_1^4 |x-1| dx \\ &= \int_0^1 (1-x) dx + \int_1^4 (x-1) dx \\ &= \left[x - \frac{x^2}{2} \right]_0^1 + \left[\frac{x^2}{2} - x \right]_1^4 \\ &= \left(1 - \frac{1}{2} \right) - 0 + (8 - 4) - \left(\frac{1}{2} - 1 \right) \\ &= 5. \end{aligned}$$

29. $ydx + (x - y^2)dy = 0$

Reducing the given differential equation to the form $\frac{dx}{dy} + Px = Q$ we get, $\frac{dx}{dy} + \frac{x}{y} = y$

That means, $P = \frac{1}{y}$, $Q = y$.

$$\therefore \text{I.F.} = e^{\int P dy} = e^{\int \frac{1}{y} dy} = e^{\log y} = y.$$

The general solution is given by $x(\text{I.F.}) = \int Q \times (\text{I.F.}) dy$

$$\Rightarrow xy = \int y^2 dy$$

$$\Rightarrow xy = \frac{y^3}{3} + C, \text{ which is the required general solution.}$$

OR

$$xdy - ydx = \sqrt{x^2 + y^2} dx$$

It is a Homogeneous Differential Equation as, $\frac{dy}{dx} = \frac{\sqrt{x^2 + y^2} + y}{x}$

$$\Rightarrow \frac{dy}{dx} = \sqrt{\frac{x^2 + y^2}{x^2}} + \frac{y}{x}$$

$$\Rightarrow \frac{dy}{dx} = \sqrt{1 + \left(\frac{y}{x}\right)^2} + \frac{y}{x} \text{ i.e., } \frac{dy}{dx} = f\left(\frac{y}{x}\right).$$

To solve, put $y = vx \Rightarrow \frac{dy}{dx} = v + x \frac{dv}{dx}$

So, $v + x \frac{dv}{dx} = \sqrt{1 + v^2} + v$

Separating variables, we get $\frac{dv}{\sqrt{1 + v^2}} = \frac{dx}{x}$

Integrating, we get $\log|v + \sqrt{1 + v^2}| = \log|x| + \log K, K > 0$

$$\Rightarrow \log|y + \sqrt{x^2 + y^2}| = 2 \log|x| + \log K$$


$$\Rightarrow \log|y + \sqrt{x^2 + y^2}| = \log(x^2 K)$$

$$\Rightarrow y + \sqrt{x^2 + y^2} = \pm Kx^2$$

$$\Rightarrow y + \sqrt{x^2 + y^2} = Cx^2, \text{ which is the required general solution.}$$

30. Consider the graph shown here.

The corner points of the feasible region are C(40, 0), D(20, 0), B(40, 160), A(20, 180).



Corner point	Value of Z
D(20, 0)	8000
C(40, 0)	16000
B(40, 160)	64000 ← Max.
A(20, 180)	62000

Maximum value of Z occurs at $x = 40, y = 160$.

And, $Z_{\max} = 64000$.

31. $\int \frac{(x^3 + x + 1)}{(x^2 - 1)} dx = \int \left(x + \frac{2x + 1}{(x - 1)(x + 1)} \right) dx$

Consider $\frac{2x+1}{(x-1)(x+1)} = \frac{A}{x-1} + \frac{B}{x+1}$

$$\Rightarrow 2x+1 = A(x+1) + B(x-1)$$

On comparing the coefficients of x and constant terms on both sides, we get

$$A+B=2, A-B=1$$


On solving these equations, we get $A = \frac{3}{2}, B = \frac{1}{2}$.

$$\begin{aligned} \text{Hence, } \int \frac{(x^3+x+1)}{(x^2-1)} dx &= \int \left(x + \frac{3}{2(x-1)} + \frac{1}{2(x+1)} \right) dx \\ &= \frac{x^2}{2} + \frac{3}{2} \log|x-1| + \frac{1}{2} \log|x+1| + C \\ &= \frac{x^2}{2} + \frac{1}{2} \log|(x-1)^3| + \frac{1}{2} \log|x+1| + C \\ &= \frac{x^2}{2} + \frac{1}{2} \log|(x-1)^3(x+1)| + C. \end{aligned}$$

SECTION D

32. Consider $y = x^2 \dots (i)$ and $y = x \dots (ii)$

On solving (i) and (ii), we get the points of intersection as $(0, 0)$ and $(1, 1)$.



$$\text{Required Area} = \int_0^1 y_{\text{parabola}} dx + \int_1^2 y_{\text{line}} dx$$

$$\text{So, required area} = \int_0^1 x^2 dx + \int_1^2 x dx$$

$$\Rightarrow \text{required area} = \left[\frac{x^3}{3} \right]_0^1 + \left[\frac{x^2}{2} \right]_1^2$$

$$\Rightarrow \text{required area} = \frac{1}{3} + \frac{3}{2} = \frac{11}{6} \text{ Sq. units.}$$

33. Let $(a, b) \in N \times N$. Then we have

$ab = ba$ (by commutative property of multiplication of natural numbers)

$$\Rightarrow (a, b)R(a, b)$$

Hence, R is reflexive.

Let $(a, b), (c, d) \in N \times N$ such that $(a, b)R(c, d)$. Then $ad = bc$

$$\Rightarrow cb = da \text{ (by commutative property of multiplication of natural numbers)}$$

$$\Rightarrow (c, d)R(a, b)$$

Hence, R is symmetric.

Let $(a, b), (c, d), (e, f) \in N \times N$ such that

$$(a, b)R(c, d) \text{ and } (c, d)R(e, f).$$

Then $ad = bc, cf = de$

$$\Rightarrow adcf = bcde$$

$$\Rightarrow af = be$$

$$\Rightarrow (a, b)R(e, f)$$

Hence, R is transitive.

Since, R is reflexive, symmetric and transitive so, R is an equivalence relation on $N \times N$.

OR

Let $A \in P(X)$. Then $A \subset A$

$$\Rightarrow (A, A) \in R$$

Hence, R is reflexive.

Let $A, B, C \in P(X)$ such that

$$(A, B), (B, C) \in R$$

$$\Rightarrow A \subset B, B \subset C$$

$$\Rightarrow A \subset C$$

$$\Rightarrow (A, C) \in R$$

Hence, R is transitive.

Let $\phi, X \in P(X)$ such that $\phi \subset X$. Hence, $(\phi, X) \in R$.

But, $X \not\subset \phi$, which implies that $(X, \phi) \notin R$.

Thus, R is not symmetric.

34. The given lines are non-parallel lines. There is a unique line-segment PQ (P lying on one and Q on the other, which is at right angles to both the lines. PQ is the shortest distance between the lines. Hence, the shortest possible distance between the insects = PQ).

The position vector of P lying on the line $\vec{r} = 6\hat{i} + 2\hat{j} + 2\hat{k} + \lambda(\hat{i} - 2\hat{j} + 2\hat{k})$ is $(6 + \lambda)\hat{i} + (2 - 2\lambda)\hat{j} + (2 + 2\lambda)\hat{k}$ for some λ .

The position vector of Q lying on the line $\vec{r} = -4\hat{i} - \hat{k} + \mu(3\hat{i} - 2\hat{j} - 2\hat{k})$ is $(-4 + 3\mu)\hat{i} + (-2\mu)\hat{j} + (-1 - 2\mu)\hat{k}$ for some μ .

$$\text{Now } \overrightarrow{PQ} = (-10 + 3\mu - \lambda)\hat{i} + (-2\mu - 2 + 2\lambda)\hat{j} + (-3 - 2\mu - 2\lambda)\hat{k}$$

Since, PQ is perpendicular to both the lines

$$(-10 + 3\mu - \lambda)(1) + (-2\mu - 2 + 2\lambda)(-2) + (-3 - 2\mu - 2\lambda)(2) = 0,$$

$$\text{i.e., } \mu - 3\lambda = 4 \quad \dots(i)$$

$$\text{And } (-10 + 3\mu - \lambda)(3) + (-2\mu - 2 + 2\lambda)(-2) + (-3 - 2\mu - 2\lambda)(-2) = 0,$$

$$\text{i.e., } 17\mu - 3\lambda = 20 \quad \dots(ii)$$

Solving (i) and (ii) for λ and μ , we get $\mu = 1, \lambda = -1$.

The position vector of the points, at which they should be so that the distance between them is the shortest, are $5\hat{i} + 4\hat{j}$ and $-\hat{i} - 2\hat{j} - 3\hat{k}$

$$\text{So, } \overrightarrow{PQ} = -6\hat{i} - 6\hat{j} - 3\hat{k}$$

$$\text{The shortest distance} = |\overrightarrow{PQ}| = \sqrt{6^2 + 6^2 + 3^2} = 9 \text{ units.}$$

OR

Eliminating t between the equations, we obtain the equation of the path $\frac{x}{2} = \frac{y}{-4} = \frac{z}{4}$, which is the equation of the line passing through the origin having direction ratios 2, -4, 4. This line is the path of the rocket.

When $t = 10$ seconds, $x = 2 \times 10 = 20$, $y = -4 \times 10 = -40$, $z = 4 \times 10 = 40$

So, the rocket will be at the point $(20, -40, 40)$. Let $P(20, -40, 40)$.

$$\therefore \text{The required distance from the origin at 10 seconds, } OP = \sqrt{20^2 + 40^2 + 40^2} \text{ km} = 60 \text{ km.}$$

Now, as the distance of a given point P(\vec{p}) from a Line $\vec{r} = \vec{a} + \lambda\vec{b}$ is $\frac{|(\vec{p} - \vec{a}) \times \vec{b}|}{|\vec{b}|}$.

Here $\vec{p} = 20\hat{i} - 40\hat{j} + 40\hat{k}$, $\vec{a} = 20\hat{i} - 10\hat{j} + 40\hat{k}$ $\therefore (\vec{p} - \vec{a}) = -30\hat{j}$.

So, the distance of the point P(20, -40, 40) from $\vec{r} = 20\hat{i} - 10\hat{j} + 40\hat{k} + \lambda(10\hat{i} - 20\hat{j} + 10\hat{k})$ is

$$= \frac{|-30\hat{j} \times (10\hat{i} - 20\hat{j} + 10\hat{k})|}{|10\hat{i} - 20\hat{j} + 10\hat{k}|} \text{ km} = \frac{|-300\hat{i} + 300\hat{k}|}{\sqrt{100 + 400 + 100}} \text{ km}$$

$$= \frac{300\sqrt{2}}{10\sqrt{6}} \text{ km} = 10\sqrt{3} \text{ km}.$$

35. For $A = \begin{bmatrix} 2 & -3 & 5 \\ 3 & 2 & -4 \\ 1 & 1 & -2 \end{bmatrix}$, $|A| = 2(0) + 3(-2) + 5(1) = -1$.

Also $\text{adj } A = \begin{bmatrix} 0 & -1 & 2 \\ 2 & -9 & 23 \\ 1 & -5 & 13 \end{bmatrix}$

$$\therefore A^{-1} = \frac{\text{adj } A}{|A|} = \frac{1}{(-1)} \begin{bmatrix} 0 & -1 & 2 \\ 2 & -9 & 23 \\ 1 & -5 & 13 \end{bmatrix} = \begin{bmatrix} 0 & 1 & -2 \\ -2 & 9 & -23 \\ -1 & 5 & -13 \end{bmatrix}$$

Given equations are $2x - 3y + 5z = 11$, $3x + 2y - 4z = -5$, $x + y - 2z = -3$

Let $A = \begin{bmatrix} 2 & -3 & 5 \\ 3 & 2 & -4 \\ 1 & 1 & -2 \end{bmatrix}$, $X = \begin{bmatrix} x \\ y \\ z \end{bmatrix}$, $B = \begin{bmatrix} 11 \\ -5 \\ -3 \end{bmatrix}$.

$$\therefore AX = AB \Rightarrow X = A^{-1}B$$

$$\therefore \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 0 & 1 & -2 \\ -2 & 9 & -23 \\ -1 & 5 & -13 \end{bmatrix} \begin{bmatrix} 11 \\ -5 \\ -3 \end{bmatrix}$$

$$\Rightarrow \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 0 - 5 + 6 \\ -22 - 45 + 69 \\ -11 - 25 + 39 \end{bmatrix}$$

$$\Rightarrow \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}$$

$$\therefore x = 1, y = 2, z = 3.$$

SECTION E

36. (i) $f(x) = -0.1x^2 + mx + 98.6$, being a polynomial function, is differentiable everywhere, hence, f is differentiable in $(0, 12)$.

(ii) $f'(x) = -0.2x + m$

Since, 6 is the critical point, $f'(6) = 0$

$$\Rightarrow -0.2 \times 6 + m = 0$$

$$\Rightarrow m = 1.2.$$

(iii) $f(x) = -0.1x^2 + 1.2x + 98.6$

$$\Rightarrow f'(x) = -0.2x + 1.2 = -0.2(x - 6)$$

For $f'(x) = 0$, $-0.2(x - 6) = 0 \Rightarrow x = 6$.

In the interval	Sign of $f'(x)$ is	Conclusion
$(0, 6)$	Positive	f is strictly increasing in $(0, 6)$
$(6, 12)$	Negative	f is strictly decreasing in $(6, 12)$

OR

(iii) $f(x) = -0.1x^2 + 1.2x + 98.6$

$$\Rightarrow f'(x) = -0.2x + 1.2 \text{ and } f''(x) = -0.2.$$

For $f'(x) = 0$, $-0.2x + 1.2 = 0 \therefore x = 6$

As $f''(6) = -0.2 < 0$.

Hence, by second derivative test we can see that, $x = 6$ is a point of local maximum.


Also, the local maximum value $= f(6) = -0.1 \times 6^2 + 1.2 \times 6 + 98.6 = 102.2$.

We have $f(0) = 98.6$, $f(6) = 102.2$, $f(12) = 98.6$.

Now $x = 6$ is the point of absolute maximum and the absolute maximum value of the function $= 102.2$.

Also $x = 0, 12$ both are the points of absolute minimum and the absolute minimum value of the function $= 98.6$.

37. (i) Let $(x, y) = \left(x, \frac{b}{a}\sqrt{a^2 - x^2}\right)$ be the upper right vertex of the rectangle.



The area function, $A = 2x \times 2 \frac{b}{a} \sqrt{a^2 - x^2}$

$$\Rightarrow A = \frac{4b}{a} x \sqrt{a^2 - x^2}, \quad x \in (0, a).$$

(ii) $\frac{dA}{dx} = \frac{4b}{a} \left[x \times \frac{-x}{\sqrt{a^2 - x^2}} + \sqrt{a^2 - x^2} \right]$

$$\Rightarrow \frac{dA}{dx} = \frac{4b}{a} \times \frac{a^2 - 2x^2}{\sqrt{a^2 - x^2}} = -\frac{4b}{a} \times \frac{(\sqrt{2}x + a)(\sqrt{2}x - a)}{\sqrt{a^2 - x^2}}$$

$$\text{For } \frac{dA}{dx} = 0 \Rightarrow x = \frac{a}{\sqrt{2}}$$

So, $x = \frac{a}{\sqrt{2}}$ is the critical point.

(iii) For the values of x less than $\frac{a}{\sqrt{2}}$ and close to $\frac{a}{\sqrt{2}}$, $\frac{dA}{dx} > 0$ and for the values of x greater than $\frac{a}{\sqrt{2}}$ and close to $\frac{a}{\sqrt{2}}$, $\frac{dA}{dx} < 0$.

Hence, by the first derivative test, there is a local maximum at the critical point $x = \frac{a}{\sqrt{2}}$. Since there is only one critical point, therefore, the area of the soccer field is maximum at this critical point $x = \frac{a}{\sqrt{2}}$.

\therefore For maximum area of the soccer field, its length should be $a\sqrt{2}$ units and its width should be $b\sqrt{2}$ units.

OR

$$(iii) A = 2x \times 2 \frac{b}{a} \sqrt{a^2 - x^2}, x \in (0, a).$$

Squaring both sides, we get $Z = A^2 = \frac{16b^2}{a^2} x^2 (a^2 - x^2) = \frac{16b^2}{a^2} (x^2 a^2 - x^4), x \in (0, a)$.

(\because A is maximum when Z is maximum.)

$$\text{Now } \frac{dZ}{dx} = \frac{16b^2}{a^2} (2xa^2 - 4x^3) = \frac{32b^2}{a^2} x(a + \sqrt{2}x)(a - \sqrt{2}x)$$

$$\text{For } \frac{dZ}{dx} = 0 \Rightarrow x = \frac{a}{\sqrt{2}}$$

$$\text{Also, } \frac{d^2Z}{dx^2} = \frac{32b^2}{a^2} (a^2 - 6x^2)$$

$$\text{As } \left(\frac{d^2Z}{dx^2} \right)_{x=\frac{a}{\sqrt{2}}} = \frac{32b^2}{a^2} (a^2 - 3a^2) = -64b^2 < 0.$$

So, by the second derivative test, there is a local maximum value of Z at the critical point $x = \frac{a}{\sqrt{2}}$.

Since there is only one critical point therefore, Z is maximum at $x = \frac{a}{\sqrt{2}}$.

Hence, A is maximum at $x = \frac{a}{\sqrt{2}}$.

\therefore For maximum area of the soccer field, its length should be $a\sqrt{2}$ units and its width should be $b\sqrt{2}$ units.

38. (i) Let X be the event that the shell fired from A hits the plane and Y be the event that the shell fired from B hits the plane. The following four hypotheses are possible before the trial, with the guns operating independently:

$$E_1 = XY, E_2 = \bar{X}\bar{Y}, E_3 = \bar{X}Y, E_4 = X\bar{Y}.$$

Let E = the shell fired from exactly one of them hits the plane.

So, $P(E_1) = 0.3 \times 0.2 = 0.06$, $P(E_2) = 0.7 \times 0.8 = 0.56$, $P(E_3) = 0.7 \times 0.2 = 0.14$,
 $P(E_4) = 0.3 \times 0.8 = 0.24$; $P(E | E_1) = 0$, $P(E | E_2) = 0$, $P(E | E_3) = 1$, $P(E | E_4) = 1$.
 Now $P(E) = P(E_1).P(E | E_1) + P(E_2).P(E | E_2) + P(E_3).P(E | E_3) + P(E_4).P(E | E_4)$
 $\therefore P(E) = 0.14 + 0.24 = 0.38$.

(ii) By Bayes' theorem,

$$P(E_3 | E) = \frac{P(E_3).P(E | E_3)}{P(E_1).P(E | E_1) + P(E_2).P(E | E_2) + P(E_3).P(E | E_3) + P(E_4).P(E | E_4)}$$

$$\Rightarrow P(E_3 | E) = \frac{0.14}{0.38} = \frac{7}{19}.$$

NOTE: The four hypotheses form the partition of the sample space and it can be seen that the sum of their probabilities is 1. The hypothesis E_1 and E_2 are actually eliminated as

$$P(E | E_1) = P(E | E_2) = 0.$$

Alternative way of writing the solution:

(i) P (shell fired from exactly one of them hits the plane)

$$= P \left[\begin{array}{l} \text{(Shell from A hits the plane and Shell from B does not hit the plane)} \\ \text{or (Shell from A does not hit the plane and Shell from B hits the plane)} \end{array} \right]$$

$$= 0.3 \times 0.8 + 0.7 \times 0.2 = 0.38.$$

(ii) P (Shell fired from B hits the plane | Exactly one of them hits the plane)

$$= \frac{P(\text{Shell fired from B hits the plane} \cap \text{Exactly one of them hits the plane})}{P(\text{Exactly one of them hits the plane})}$$

$$= \frac{P(\text{Shell from only B hits the plane})}{P(\text{Exactly one of them hits the plane})}$$

$$= \frac{0.14}{0.38} = \frac{7}{19}.$$

This paper has been issued by CBSE for 2022-23 Board Exams of class 12 Mathematics (041).

Note : We've re-typed the official sample paper and, also done the necessary corrections at some places. Apart from that, further illustrations have been added as well in some questions.

If you notice any error which could have gone un-noticed, please do inform us via **WhatsApp @ +919650350480 (message only)** or, via **Email at iMathematicia@gmail.com**

Let's learn **Math** with **smile:-)**

- **O.P. GUPTA, Math Mentor**

For Detailed Solutions, please visit **[YouTube.com/MathematiciaByOPGupta](https://www.youtube.com/MathematiciaByOPGupta)**

Detailed Solutions (PTS-10)

SECTION A

01. (c) Since $P_{3 \times 2} A_{m \times n} = Q_{3 \times 2}$ implies, $2 = m$ and $n = 2$.

Therefore, the order of A must be 2×2 .

02. (b) Since $A(\text{adj.}A) = \begin{bmatrix} -2 & 0 & 0 \\ 0 & -2 & 0 \\ 0 & 0 & -2 \end{bmatrix} = -2 \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} = -2I_3$

Also, $A(\text{adj.}A) = |A| I_3$

$$\therefore |A| = -2.$$

03. (c) As \vec{a} and \vec{b} are collinear so, $\frac{\alpha}{2} = \frac{3}{-1} = \frac{-6}{-\beta}$

$$\text{Considering } \frac{\alpha}{2} = \frac{3}{-1}, \frac{3}{-1} = \frac{-6}{-\beta}$$

$$\Rightarrow \alpha = -6, \beta = -2$$

$$\therefore (\alpha + \beta) = -8.$$

04. (c) As the greatest integer function is not differentiable at integral points. So, here $f(x)$ will be non-differentiable at $x = 1$.

05. (b) $\int e^x \left(\log \sqrt{x} + \frac{1}{2x} \right) dx = e^x \times \log \sqrt{x} + C.$

Using $\int e^x [f(x) + f'(x)] dx = e^x f(x) + C$, where $f(x) = \log \sqrt{x}$, $f'(x) = \frac{1}{2x}$.

06. (a) As there are no arbitrary constants in the particular solution of a differential equation so, the number of arbitrary constants is 0.

07. (b) a linear function to be optimized.

08. (c) Required length of perpendicular drawn from $(4, -7, 3)$ on y -axis $= \sqrt{4^2 + 3^2} = 5$ units.

09. (c) $\int_1^e \frac{\log x}{x} dx = \frac{1}{2} [(\log x)^2]_1^e = \frac{1}{2} [(\log e)^2 - (\log 1)^2]$
 $= \frac{1}{2} [1 - 0] = \frac{1}{2}.$

10. (d) $\begin{vmatrix} 2 & 3 & 2 \\ x & x & x \\ 4 & 9 & 1 \end{vmatrix} + 3 = 0$

On expanding along R_1 , we get : $2(x - 9x) - 3(x - 4x) + 2(9x - 4x) + 3 = 0$

$$\Rightarrow 2(-8x) - 3(-3x) + 2(5x) + 3 = 0$$

$$\Rightarrow -16x + 9x + 10x + 3 = 0$$

$$\Rightarrow 3x + 3 = 0$$

$$\therefore x = -1.$$

11. (a) As maximum value of z occurs at $(2, 4)$ and $(4, 0)$ so, $a(2) + b(4) = a(4) + b(0)$
 $\Rightarrow a = 2b.$

12. (c) As $\begin{bmatrix} 2 & -1 & 3 \\ \lambda & 0 & 7 \\ -1 & 1 & 4 \end{bmatrix}$ is non-invertible so, $\begin{vmatrix} 2 & -1 & 3 \\ \lambda & 0 & 7 \\ -1 & 1 & 4 \end{vmatrix} = 0$

$$\Rightarrow 2(-7) + 1(4\lambda + 7) + 3(\lambda) = 0 \quad \therefore \lambda = 1.$$

13. (a) $a_{12} + a_{22} = |1^2 - 2| + |2^2 - 2| = 1 + 2 = 3.$

14. (b) Re-writing the D.E., $\frac{dy}{dx} + \frac{y}{\sqrt{x}} = \frac{e^{-2\sqrt{x}}}{\sqrt{x}}$

On comparing to $\frac{dy}{dx} + Py = Q$, we observe $P = \frac{1}{\sqrt{x}}$ and $Q = \frac{e^{-2\sqrt{x}}}{\sqrt{x}}$

\therefore Integration factor $= e^{\int \frac{1}{\sqrt{x}} dx} = e^{2\sqrt{x}}.$

15. (c) $P(B' | A) = \frac{P(B' \cap A)}{P(A)}$

$\Rightarrow P(B' | A) = \frac{P(A) - P(A \cap B)}{P(A)} \quad (\because B' \cap A = A - B)$

$\Rightarrow P(B' | A) = \frac{P(A) - P(A) \times P(B)}{P(A)} \quad (\because A \text{ and } B \text{ are independent events})$

$\Rightarrow P(B' | A) = 1 - P(B)$

$\Rightarrow P(B' | A) = 1 - \frac{1}{4} = \frac{3}{4}.$

16. (c) $f(x)$ is discontinuous at exactly three points, $x = 0, 1, -1.$

17. (d) As the number of Reflexive relations defined on a set of n elements $= 2^{n(n-1)}.$
So, 2^6 reflexive relations are possible in the set A where $n(A) = 3.$

18. (b) Equation of line joining $(-1, 3, 2)$ and $(5, 0, 6)$ is $\frac{x+1}{6} = \frac{y-3}{-3} = \frac{z-2}{4} = \lambda$

The random point on the line is $(6\lambda - 1, -3\lambda + 3, 4\lambda + 2).$

As x -coordinate of point P is 2 so, $6\lambda - 1 = 2 \Rightarrow \lambda = \frac{1}{2}$

Therefore, the z -coordinate is $4\lambda + 2 = 4\left(\frac{1}{2}\right) + 2 = 4.$

19. (b) Here both A and R are true and R is not the correct explanation of $A.$

20. (c) $\overrightarrow{OP} = \frac{2\overrightarrow{OB} + 1\overrightarrow{OA}}{2+1}$

$\Rightarrow \overrightarrow{OP} = \frac{2(2\hat{i} - \hat{j} + 2\hat{k}) + 1(2\hat{i} - \hat{j} - \hat{k})}{3}$

$\therefore \overrightarrow{OP} = 2\hat{i} - \hat{j} + \hat{k}.$ So, A is true.

Also, R is false. Since $\vec{a} \cdot \vec{b} = |\vec{a}| |\vec{b}| \cos \theta.$

SECTION B

21. $\sin^{-1} \left[\sin \left(-\frac{17\pi}{8} \right) \right] = \sin^{-1} \left[\sin \left(-2\pi - \frac{\pi}{8} \right) \right] = \sin^{-1} \left[-\sin \left(\frac{\pi}{8} \right) \right] = -\sin^{-1} \left(\sin \frac{\pi}{8} \right) = -\frac{\pi}{8}.$

OR

Note that the domain of $\sin^{-1} x$ is $x \in [-1, 1]$ and that of $\tan^{-1} x$ is $x \in \mathbb{R}.$

So, the domain of $f(x) = \tan^{-1} x + \frac{1}{2} \sin^{-1} x$ is $x \in [-1, 1].$

Answers (PTS-16)

- 01.** (d) **02.** (d) **03.** (c) **04.** (b) **05.** (c) **06.** (c) **07.** (b)
08. (b) **09.** (d) **10.** (b) **11.** (c) **12.** (c) **13.** (b) **14.** (d)
15. (d) **16.** (c) **17.** (d) **18.** (d) **19.** (d) **20.** (a)
- 21.** $2 \sin^{-1} x$ **22.** $0.32 \pi \text{ cm}^2/\text{s}$ **23.** $2\hat{i} - \hat{j} + \hat{k}$ **24.** **OR** $X = \begin{pmatrix} 1 & -2 \\ 2 & 0 \end{pmatrix}$
- 25.** $\sin^{-1} \left(\frac{e^x + 2}{3} \right) + C$ **OR** $x \cos 2a - \sin 2a \log |\sin(x+a)| + C$ **26.** f is not onto.
- 27.** $f(x)$ is increasing on $x \in [0, 2]$ and decreasing on $x \in (-\infty, 0] \cup [2, \infty)$
- 28.** $y = x \log \left| \frac{x}{(x-y)^2} \right|$ **OR** $xe^x - e^x + 1 = \sqrt{1-y^2}$ **29.** $\mu = \frac{1}{4}$
- 30.**

X	0	1	2
P(X)	$\frac{9}{87}$	$\frac{40}{87}$	$\frac{38}{87}$


Mean = $\frac{116}{87}$
- 31.** **OR** $f(x)$ is not differentiable at $x=1$, $f(x)$ is differentiable at $x=2$
- 32.** 4π Sq. units **33.** $\left(\frac{1}{2}, -\frac{1}{2}, -\frac{3}{2} \right)$
- 34.** Maximum value of z is 10. **OR** Maximum value of z is $22\frac{8}{13}$.
- 35.** **OR** $2x^2 - 3x + 1$
- 36.** (i) $x + y + z = 7000$, $x - y = 0$, $10x + 16y + 17z = 110000$
 (ii) $\begin{pmatrix} 1 & 1 & 1 \\ 1 & -1 & 0 \\ 10 & 16 & 17 \end{pmatrix} \begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} 7000 \\ 0 \\ 110000 \end{pmatrix}$
 (iii) System of equations is consistent and, the system of equations will have unique solution as, $|A| \neq 0$. **OR** (iii) ₹1125/-, ₹4750/-.
- 37.** (i) $-4\hat{i}$ (ii) $-\frac{5}{2}\hat{i} + \frac{3\sqrt{3}}{2}\hat{j}$ (iii) $\frac{3}{2}\hat{i} + \frac{3\sqrt{3}}{2}\hat{j}$, $6\sqrt{3}\hat{k}$ **OR** (iii) $3\sqrt{3}$ Sq. units.
- 38.** (i) 0.039 (ii) $\frac{5}{13}$.

Answers (PTS-17)

- 01.** (b) **02.** (c) **03.** (d) **04.** (d) **05.** (c) **06.** (b) **07.** (a)
08. (d) **09.** (b) **10.** (d) **11.** (a) **12.** (c) **13.** (b) **14.** (d)
15. (c) **16.** (b) **17.** (d) **18.** (b) **19.** (c) **20.** (a)
- 21.** $\pi - 2 \sin^{-1} x$ or, $2 \cos^{-1} x$ (both answers are possible) **22.** $9 \text{ cm}^3/\text{s}$
- 23.** **OR** $2, -3, 0$; $\vec{r} = -3\hat{i} + 5\hat{j} - 2\hat{k} + \lambda(2\hat{i} - 3\hat{j})$ **24.** $\frac{y \sin(xy)}{\sin 2y - x \sin(xy)}$
- 25.** $\frac{1}{\sqrt{14}}, \frac{2}{\sqrt{14}}, \frac{3}{\sqrt{14}}$

Reviews for MATHMISSION by O.P. GUPTA

Class 12 Books (Term I & II) for Maths (041)



Amazon Customer

★★★★★

Excellent Book from an excellent Teacher

Reviewed in India on 13 September 2021

Thank you so much sir for this wonderful book. I appreciate your effort and I am sure that this will be an excellent reference book for both teachers and students those who are going to appear for coming board examination. Here we will get questions of all levels which helped me a lot in preparing question papers for my term 1 revision exams. I recommend this to all students and teachers

Manish Saxena

★★★★★

BEST BOOK

Reviewed in India on 11 September 2021

Bahut achi book hai. Agar students is book se term 1 ki preparation karange to full marks ayange. Abhi tak Maine jitni book padhi usme sabse best hai.

5★ Perfect product!

Many questions included in this book more helpful for class 12 students must buy book for maths in this price loved it

Flipkart Customer, Muvattupuzha

Certified Buyer

Brij Mohan Negi

★★★★★

The best book All types of questions are covered

Reviewed in India on 11 September 2021

For class 12 students the best book

All types of questions are covered

U can score full marks by doing this book

MANDEEP SINGH

★★★★★

Excellent Book

Reviewed in India on 12 September 2021

Very good book for the preparation of Term-I CBSE

Examinations. Students can get good marks in the first

term examinations. Thanks Gupta ji.

Pardeep sharma

★★★★★

Great book

Reviewed in India on 12 September 2021

This book is, presently need of the hour! As per latest cbse pattern term 1 ! Being a mathematics teacher I recommend the same for every student going to appear in term 1 of maths exam in coming nov 2021!

Maths Teacher

★★★★★

Maths mission book class12

Reviewed in India on 11 September 2021

Best one for class 12 (MCQTERM - 1) CBSE students to buy and practice the sums.

I am sure that success is yours.

Smitha

★★★★★

Very nice book, must buy for MCQ type questions

Reviewed in India on 12 September 2021

A variety of tricky MCQs, assertion reason, case study, source based questions are available which will definitely train the students to face their TERM 1 exam with confidence

Deepika Bhati

★★★★★

Very good collection of questions..

Reviewed in India on 11 September 2021

Recommended to all students ...

Really appreciable work done by Mr. O P Gupta

Sir...Book for all type of questions from zero level to high level with complete explanations.

Sachin Pandey

★★★★★

TOTAL PACAKGE (FOR TERM 1) CBSE EXAM 12TH

Reviewed in India on 10 September 2021

BEST BOOK FOR PREPARATION. IT HAS ALL THE THINGS REQUIRED FOR PREPARATION NOT ONLY FOR TERM -1 BUT ALSO FOR OTHER EXAM. FULLY SOLVED..AND IT IS BEST BOOK IN THE MARKET

Ashita

★★★★★

MATH MISSION WILL CREATE MAGIC !!

Reviewed in India on 11 September 2021

Math mission is highly recommendable to all the students of class 12th. This book is comprising of every type of question related to CHANGED PATTERN AND SYLLABUS OF TERM 1. Most informative and helpful reference book of Mathematics till date.

Neetha

★★★★★

Latest supporting material for class 12

Reviewed in India on 12 September 2021

The book is as per the latest cbse curriculum and include all types of questions like Assertion- reasoning, source based and MCQ's of all types. Really a supporting material for students and educators

To order your copy of MATHMISSION Book of class XII, WhatsApp @ +91-9650350480

Class XII

For CBSE Exams (2022-23)

- ✓ FORMULAE CLUSTER
- ✓ SUBJECTIVE TYPE QUESTIONS
- ✓ MULTIPLE CHOICE QUESTIONS
- ✓ CASE STUDY QUESTIONS
- ✓ ASSERTION - REASONING QUESTIONS
- ✓ COMPETENCY BASED QUESTIONS



MATH MISSION

O.P. GUPTA
INDIRA AWARD WINNER

ABOUT THE AUTHOR



O.P. GUPTA, having taught Math passionately over a decade, has devoted himself to this subject. Every book, study material or practice sheets, tests he has written, tries to teach serious Math in a way that allows the students to learn Math without being afraid. His resources have helped students and teachers for a long time across the county. He has contributed in CBSE Question Bank (issued in April 2021). Mr GUPTA has been invited by many educational institutions for hosting sessions for the students of senior classes. Being qualified as an Electronics & Communications engineer, he has pursued his graduation later on with Math from University of Delhi due to his passion towards Mathematics. He has been honored with the prestigious INDIRA AWARD by the Govt. of Delhi for excellence in education.

OUR OTHER PUBLICATIONS



ONLINE SUPPORT | For CBSE Board Papers, Sample Papers, Topic Tests, NCERT Solutions & More...



www.theopgupta.com

**Do You Have Any
Queries Regarding Maths?**

Feel free to contact us

✉ iMathematicia@gmail.com

☎ +919650350480 (Message Only)



Buy our Math books at www.theopgupta.com/shop